



*12<sup>th</sup> Meeting of the  
International Scientific Committee  
for Tuna and Tuna-Like Species in the North Pacific Ocean  
Sapporo, Hokkaido, Japan  
18-23 July 2012*

## **The 2011 Canadian North Pacific Albacore Troll Fishery<sup>1</sup>**

**John Holmes**

Fisheries and Oceans Canada  
Pacific Biological Station  
3190 Hammond Bay Road  
Nanaimo, BC, Canada, V9T 6N7

July 2012

<sup>1</sup>Prepared for the Twelfth Meeting of the International Scientific committee on Tuna and Tuna-like Species in the North Pacific Ocean (ISC), 18-23 July 2012, Sapporo, Hokkaido, Japan. Document should not be cited without permission of the authors.

## 1.0 INTRODUCTION

The Canadian fishery for highly migratory species uses troll gear with jigs to target juvenile north Pacific albacore tuna (*Thunnus alalunga*) in the surface waters of the Pacific Ocean. The majority of catch and effort by the Canadian fleet has occurred in the coastal waters of Canada and the United States since 2000. However, some vessels in the fleet follow albacore tuna concentrations into offshore waters and into the central and western Pacific Ocean.

Management regulations for Canadian vessels fishing albacore tuna in 2011 are documented in the Pacific Region Integrated Fisheries Management Plan: Tuna - April 1, 2011 to March 31, 2012, which is available at <http://www.dfo-mpo.gc.ca/Library/343252.pdf>. Although these regulations specify that Canadian tuna fishery in Pacific Ocean waters was open from 01 April 1 2011 to 31 March 2012, all catch and effort occurred between 15 June and 31 October 2011. Canadian vessels are licensed to fish within the Canadian exclusive economic zone (EEZ), the EEZ of the United States, and the highseas (waters outside the EEZs) in both the north and south Pacific Oceans. Access to the United States Exclusive Economic Zone (EEZ) is permitted through a bilateral Treaty, which provides for reciprocal access by Canadian-flagged and US-flagged vessels licensed to fish for albacore and for the landing of albacore catches at designated ports within each country.

The present report summarizes Canadian Category I (total annual catch and effort, catch per unit of effort (CPUE)) for 1995-2010 period and provides provisional Category I, Category II (logbook catch and effort data summarized on 1° x 1° grid), and Category III (bycatch, catch size composition) data from the 2011 fishery. Summaries of the 1995 through 2010 fisheries are presented by Shaw (1997, 1999), Shaw and Argue (1999, 2000), Argue and Shaw (2000), Shaw and Stocker (2002), Stocker and Shaw (2004a, 2004b, 2005), Stocker (2006, 2007), and Holmes (2008, 2009, 2010, 2011).

## 2.0 DATA SOURCES

Data on albacore tuna catch and effort from 1995 through to the present are compiled from hail records, logbooks, and sales slips from processing plants and stored in the Canadian Albacore Tuna Catch and Effort Relational Database (Stocker et al. 2007). This database generates the best available estimates of total annual catch and effort by geographic area for the Canadian fishery. All Canadian fishing vessels are required to hail a third party service provider when they intend to start fishing and when fishing ceases. At present, the hail data are used primarily to ensure that fishing activity in the United States EEZ ceases by 31 October as required by the Canada-United States Albacore Tuna Treaty, but they could be used to estimate fleet size in the three fishing zones if necessary. Canadian vessels must also carry logbooks in which daily catch and effort data (latitude, longitude, number of fish, estimated weight) are recorded for albacore and non-target species. These data have the highest temporal and spatial resolution are obtained from completed copies of the logbooks submitted at the end of the fishing season. The third data source, sales slips, record the weight of albacore landed and bought by a domestic Canadian buyer and provide the most accurate estimates of albacore catch in weight since these data are the basis for payment to harvesters. In some years, sales slips underestimate total catch because

they do not fully account for landings at international ports, domestic public sales or take-home totals (Stocker et al. 2007). Logbooks, sales slips from domestic buyers, and at-sea trans-shipment slips, completed at the time fish are landed and sold, must be returned to Fisheries and Oceans Canada (DFO) for entry into the albacore catch database (Stocker et al. 2007).

Fork length data were collected from the Canadian catch landed in United States ports by the US port sampling program and forwarded to Canada prior to 2009. In 2009, Canada implemented an on-board harvester sampling program to obtain fork lengths of fish in the Canadian catch. Participation is voluntary and harvesters are asked to measure first 10 albacore landed on a daily basis and record these data in the spaces provided in their logbooks. Fork length data reported by Canada since 2009 (Holmes 2010, 2011, present report) are from the domestic on-board sampling program only.

The preliminary catch and effort data for 2011 in this report were taken from database version 12.04.23. Logbook coverage was estimated to be 98% of expanded catch and effort. The data in this report up to and including 2010 are considered definitive and are derived from a reconciliation of trip log (best estimates of effort, catch in pieces, and geographic location) and sales slip (best estimate of catch weight) data (Stocker et al. 2007). In some cases, Category I data from the 1995-2010 period have been revised relative to previous submissions by Canada due to the receipt of outstanding logbooks and sales slip reconciliation update of catch weights. See Holmes (2012) for details.

### **3.0 CATEGORY I DATA**

#### **3.1 Catch**

The preliminary estimate of the Canadian catch of north Pacific albacore in 2011 is 5,393 metric tons (t) and is an 18% decrease relative to 2010 (Table 1; Figure 1). The total catch of north Pacific albacore tuna by the Canadian troll fishery has ranged from 1,761 t in 1995 to 7,857 t in 2004 and averaged  $4,857 \pm 1,753$  (sd) t for the 1995 to 2010 period. All of the 2011 catch except for 5 t occurred in FAO Statistical Area 67 (Table 2), continuing a trend of concentrating catch and effort in eastern Pacific Ocean (EPO) that began in 2005. The 2011 catch was distributed among the Canadian EEZ (12.5%), the United States EEZ (86%), and highseas waters (1.5%). The Canadian fishery operated north of the equator primarily within the IATTC convention area, but a minor amount of catch (1 t) was made in the WCPFC convention area. No Canadian vessels were active in the south Pacific Ocean in 2011.

#### **3.2 Effort**

The Canadian albacore troll fleet consisted of 177 unique vessels in 2011, which is an 11% increase in fleet size relative to 2010, but below the average of 212 for the 1995-2010 period (Table 1). All Canadian troll vessels targeted the north Pacific albacore stock exclusively.

Fishing effort in the Canadian tuna fishery is measured as the number of vessel fishing days (v-d). The 2010 estimate of fishing effort is 8,568 v-d and is a 13 % increase in effort relative to

effort in 2010 (Table 1; Figure 1). Annual fishing effort has ranged between 4,320 v-d in 1997 and 10,021 v-d in 2001, averaging  $7,490 \pm 1,551$  (sd) v-d since 1995.

#### 4.0 CATEGORY II DATA

The Canadian troll fleet operated between 38 and 54 °N latitude and from the west coast of North America to 151° W in 2011 (Figures 2 and 3), which is further south and north than in the previous five years, but consistent with the pattern established since 2006 of fishing primarily within the Inter-American Tropical Tuna Commission (IATTC) convention area east of 150 °W. Approximately 76% of the fishing effort occurred within the US EEZ and resulted in 86% of the total catch in 2011, similar to average proportions of effort and catch of 78% and 79%, respectively, for the 2000-2009 period (period for which logbook coverage exceeds 80% of the catch annually). In contrast, 21% of the effort occurred in Canadian waters, resulting in only 13% of the catch and 2% of the catch and 3% of the effort occurred in adjacent highseas waters. The highest catch and effort occurred within the coastal waters of Canada and the United States, particularly along the Oregon coastline (Figures 2 and 3).

The seasonal pattern of nominal catch-rates in 2011 for the Canadian fishery departs substantially from the average seasonal pattern for 2000-2009 (Figure 4). On average, catch-rate increases to a peak in July-August and then declines to low values by late October. In contrast, nominal 2011 catch-rates exhibit a minor peak in early July followed by a steady increase from late August to the end of October, with the highest catch-rates occurring at the end of the fishing season (Figure 4).

Albacore were caught in waters with sea surface temperatures ranging between 10 and 20 °C in 2010, but the most productive waters were between 14 and 18 °C and accounted for 99% of the catch (Figure 5). In contrast, the majority of catch has occurred in waters between 16 and 18 °C in other years.

#### 5.0 CATEGORY III DATA

##### 5.1 Bycatch

Reported bycatch in 2011 consisted of 71 fish and eight species (Table 3). Ten albacore considered to small to retain were recorded in June and July, 8 blue sharks, 2 shortfin mako sharks, 6 Pacific bluefin tuna, 1 bigeye tuna, 2 mahi mahi, and 42 yellowtail *Seriola lalandi* were captured and released alive. Total weight of all bycatch is estimated to be approximately 951 kg, of which the shortfin mako sharks accounted for 300 kg.

##### 5.2 Biological

Forty-three (43) vessels measured fork lengths and compiled catch size frequency data in 2011. A total of 14,338 fork length measurements were recorded (Figure 6), representing a sample of 1.72% of the total reported catch (831,299 fish). This sampling rate is well above the target rate of 1.0% of the reported Canadian catch. The majority of measurements (79%) were made on albacore caught in the US EEZ, where the majority of Canadian fishery effort occurred.

Albacore in the Canadian catch ranged from 50 cm to 90 cm in size (Figure 6) and overall a single mode at 65-68 cm was dominant in the pooled data, corresponding to 2-yr old fish. However, categorizing the data by fishing area revealed two modes at 67 and 76 cm (2- and 3-yr old fish) among fish sampled in the Canadian EEZ but only a single mode at 65-68 cm among fish sampled in the US EEZ and the highseas (Figure 6). The most common pattern of length-frequency data sampled from the Canadian catch is a single mode between 64 and 68 cm FL, corresponding to 2-year old fish. A two-mode pattern is rarer, but has been observed several times and usually occurs when albacore shift their distribution in coastal waters northward in response to ocean conditions.

## 6.0 DISCUSSION

The 2011 Canadian albacore fishery was remarkable for the fact that albacore availability in coastal waters was delayed until late in the fishing season (September-October) and the highest abundance occurred with 2-3° of longitude of the Oregon and Washington coastlines (Figure 2). This reduced availability is reflected by a reduction in catch and increase in effort in 2011 relative to the 2006-2010 period, in which catch was relatively stable and effort was declining (Figure 1). The highest catches of albacore occurred in cooler waters (14-18 °C) than usual (Figure 5) and the size frequency data reveal that albacore caught in the Canadian EEZ consisted of two age groups, rather than the single age-group in other areas (Figure 6). The larger age-3 fish commonly enter the Canadian catch if substantial effort occurs south of 40°N or the juvenile albacore shift north due to ocean conditions as occurred in 2010. The size frequency pattern in the Canadian EEZ is consistent with older and larger fish migrating further north than younger fish, but the explanation for absence of a second mode in the US EEZ size frequency data is not clear at present.

## 7.0 RESEARCH

Canada is beginning a tagging program using pop-up satellite tags to investigate the movement patterns of juvenile albacore in the eastern Pacific Ocean. This project is responding to a need to understand the movement patterns of albacore and bring movement into the stock assessment model that was identified by the Albacore Working Group during the 2011 stock assessment workshop. The goal is to deploy up to 50 tags on juvenile albacore in the eastern Pacific Ocean in a least two locations (near 150° W, coastal waters of North America, approximately 125-130° W) and two times (early spring at the western location, late summer or fall in the coastal location). Pop-up satellite tags are planned for this project in an attempt to ensure that biases associated with unexpectedly low returns that have affected interpretation of other albacore tagging programs, particularly in the western Pacific Ocean. Current effort is focused on developing handling and tagging protocols and minimizing premature tag release from fish.

Canada is also conducting research evaluating the impacts of biological and oceanographic variables on the population dynamics of albacore tuna in response to a need identified by the Albacore Working Group during the 2011 stock assessment workshop. A Bayesian production model previously presented to the Working Group is the basis of this research. Conventional

biomass dynamics model assume that the carrying capacity (K) and the intrinsic rate of population growth (r) of a stock are temporally invariant and usually estimate these parameters based on catch rates (such as CPUE) and fishing effects from a single fishery. The model developed for this research uses data from multiple fisheries and gears and allows K and r to change over the years. As a first step, the effects of oceanographic indices such as the Pacific Decadal Oscillation (PDO), North Pacific Oscillation Index (NOI), Multivariate ENSO Index (MEI), and the North Pacific Gyre Oscillation (NPGO) on K and r are modeled and the significance of each oceanographic variable is examined based on the model fitting. Based on these first order results, other more complex stock assessment models, for example an age-structured production model, would also be constructed, and oceanographic variables would then be incorporated into the models and the behaviour of alternative models will be statistically compared to produce a reasonably reliable and robust ecological model for the north Pacific albacore tuna population.

Preliminary results show the NPGO significantly affects productivity in a positive manner for a time lag up to 5 years and the probability of this impact is highest when a 4-year time lag is used. At time lags exceeding 5 years, the NPGO has no significant impact on productivity and the model performance does not statistically improve relative the base model.

## 8.0 REFERENCES

- Argue, A.W. and W. Shaw. 2000 MS. An update for Canadian tuna fisheries in the north and south Pacific ocean. 17pp. Document submitted by DFO to the Thirteenth Meeting of the Standing Committee on Tuna and Billfish in Noumea, New Caledonia, July 5-12, 2000.
- Holmes, J.A. 2008 MS. The 2007 Canadian North Pacific albacore troll fishery. Document prepared for the Eighth Meeting of the International Scientific Committee on Tuna and Tuna-like Species in the North Pacific Ocean (ISC), 22-27 July 2008, Takamatsu, Japan. ISC/08/Plenary/04: 13 p.
- Holmes, J.A. 2009 MS. The 2008 Canadian north Pacific albacore troll fishery. Document prepared for the Ninth Meeting of the International Scientific Committee on Tuna and Tuna-like Species in the North Pacific Ocean (ISC), 15-20 July 2009, Kaohsiung, Taiwan. ISC/09/Plenary/07: 16 p.
- Holmes, J.A. 2010 MS. The 2009 Canadian North Pacific albacore troll fishery. Document prepared for the Tenth Meeting of the International Scientific Committee on Tuna and Tuna-like Species in the North Pacific Ocean (ISC), 21-26 July 2010, Victoria, Canada. ISC/10/Plenary/10: 15 p.
- Holmes, J.A. 2011 MS. The 2010 Canadian North Pacific albacore troll fishery. Document prepared for the Eleventh Meeting of the International Scientific Committee on Tuna and Tuna-like Species in the North Pacific Ocean (ISC), 20-25 July 2011, San Francisco, USA. ISC/11/Plenary/08: 18 p.
- Holmes, J.A. 2012 MS. Revised Canadian albacore fishery statistics, 1995-2010, and provisional fishery statistics for the 2011 albacore troll fishery. Working paper prepared for the ISC Albacore Working Group Meeting, 14-15 July 2012, Sapporo, Japan, ISC/12/ALBWG/01: 18 p.
- Kleiber, P., and C. Perrin. 1991. Catch-per-effort and stock status in the U.S. north Pacific albacore fishery: reappraisal of both. *Fish. Bull.*, US 89: 379-386.
- Shaw, W. 1997 MS. The Canadian albacore fishery off the west coast of British Columbia, 1996-97. 15 pp. Document submitted by DFO to the Fifteenth Meeting of the North Pacific Albacore Workshop, Nanaimo, B.C., December 1997.
- Shaw, W. 1999 MS. An update of the Canadian north Pacific albacore fishery. 6 pp. Document submitted by DFO to the Second Meeting of the Interim Scientific Committee for Tuna and Tuna-Like Species in the North Pacific Ocean, Honolulu, Hawaii, January 1999.
- Shaw, W. and A.W. Argue. 1999 MS. The Canadian albacore fisheries in the north Pacific ocean, 1995-1998. Document submitted by DFO to the Sixteenth Meeting of the North Pacific Albacore Workshop, Kesenuma, Japan, November 4-6, 1999.

- Shaw, W. and A.W. Argue. 2000 MS. The 1999 Canadian north Pacific albacore troll fishery. Document submitted by DFO to the Seventeenth Meeting of the North Pacific Albacore Workshop, Taipei, Taiwan, December 6-13, 2000.
- Shaw, W. and M. Stocker. 2002 MS. The 2000 and 2001 Canadian North Pacific albacore troll fishery. Document submitted by DFO to the Eighteenth Meeting of the North Pacific Albacore Workshop, La Jolla, CA, December 4-11,2002. NPALB/02/16: 22p.
- Stocker, M. 2006 MS. The 2005 Canadian North Pacific albacore troll fishery. Document submitted by DFO to the ISC Albacore Working Group Meeting, 2006 National Research Institute of Far Seas Fisheries (NRIFSF), 5-7-1 Orido Shimizu-ku, Shizuoka-shi, 424-8633 Japan. ISC-ALBWG/06/05: 10p.
- Stocker, M. 2007 MS. The 2006 Canadian North Pacific albacore troll fishery. Document submitted by DFO to the ISC Plenary Session, Busan, Korea. ISC/07/PLENARY/04: 8p.
- Stocker, M. and W. Shaw. 2004a MS. The 2002 Canadian north Pacific albacore troll fishery. Document submitted by DFO to the Fourth Meeting of the Interim Scientific Committee on Tuna and Tuna-like Species in the North Pacific Ocean (ISC), Honolulu, Hawaii, 26 January-4 February, 2004. ISC/04/Plenary/1: 11p.
- Stocker, M. and W. Shaw. 2004b MS. The 2002 and 2003 Canadian north Pacific albacore troll fishery. Document submitted by DFO to the 19<sup>th</sup> Meeting of the North Pacific Albacore Workshop, Nanaimo, BC, November 25-December 2, 2004. NPALB/04/03: 11p.
- Stocker, M. and W. Shaw. 2005 MS. The 2004 Canadian north Pacific albacore troll fishery. Document submitted by DFO to the ISC Albacore Working Group Meeting, NOAA Fisheries, Southwest Fisheries Science Center, La Jolla, CA, USA, November 28-December 2, 2005. ISC-ALBWG/05/12: 10p.
- Stocker, M., H. Stiff, W. Shaw, and A.W. Argue. 2007. The Canadian albacore tuna catch and effort relational database. Canadian Technical Report of Fisheries and Aquatic Sciences 2701: vi+76 p.



**Table 1.** Fishery statistics from the Canadian troll fishery for north Pacific albacore tuna, 1995-2011.

<b>Year</b>	<b>Total Catch (t)</b>	<b>Effort (vessel- days)</b>	<b>Total Vessels</b>	<b>Logbook Coverage (%)</b>
1995	1,761	5,923	287	22%
1996	3,321	8,164	295	50%
1997	2,166	4,320	200	57%
1998	4,177	6,018	214	79%
1999	2,734	6,970	238	89%
2000	4,531	8,769	243	83%
2001	5,248	10,021	248	94%
2002	5,379	8,323	232	84%
2003	6,861	8,429	193	98%
2004	7,857	9,942	221	97%
2005	4,888	8,578	213	99%
2006	6,008	6,277	174	100%
2007	6,667	6,961	207	100%
2008	5,476	5,919	134	95%
2009	5,690	6,553	138	99%
2010	6,552	7,592	159	98%
2011 <sup>1</sup>	5,393	8,568	177	99%

1. 2011 data are preliminary based on Ver.12.04.23 of the *Canadian Albacore Tuna Catch and Effort Relational Database*.
2. (Reported Catch/Expanded Catch) x 100

**Table 2.** Total Canadian catch (t) of north Pacific albacore tuna in FAO Statistical Areas in the north Pacific Ocean.


Year	FAO Statistical Area			Total
	NE Pacific (67)	NW Pacific (61)	EC Pacific (77) <sup>2</sup>	
2002	5,089	152	138	5,379
2003	6,429	341	91	6,861
2004	7,710	44	102	7,856
2005	4,877	11	0	4,888
2006	6,007	0	0	6,007
2007	6,666	0	1	6,667
2008	5,476	0	0	5,476
2009	5,689	0	1	5,690
2010	6,552	0	0	6,552
2011 <sup>1</sup>	5,388	0	5	5,393

<sup>1</sup> Preliminary 2011 data obtained Version 12.04.23 of the *Canadian Albacore Tuna Catch and Effort Relational Database*.


<sup>2</sup> Excludes catch data from south of the equator.

**Table 3.** Reported catch of non-target species (by-catch) by the Canadian albacore troll fishery in 2011.


Month	Common name	Scientific Name	No. of fish
June	Albacore	<i>Thunnus alalunga</i>	7
	Mahi mahi	<i>Coryphaena hippurus</i>	2
July	Albacore	<i>Thunnus alalunga</i>	3
	Blue shark	<i>Prionace glauca</i>	3
	Pacific bluefin tuna	<i>Thunnus orientalis</i>	2
	Yellowtail amberjack	<i>Seriola lalandi</i>	8
August	Blue shark	<i>Prionace glauca</i>	3
	Pacific bluefin tuna	<i>Thunnus orientalis</i>	2
	Shortfin mako shark	<i>Isurus oxyrinchus</i>	1
	Yellowtail amberjack	<i>Seriola lalandi</i>	9
September	Bigeye tuna	<i>Thunnus obesus</i>	1
	Blue shark	<i>Prionace glauca</i>	2
	Pacific bluefin tuna	<i>Thunnus orientalis</i>	2
	Shortfin mako shark	<i>Isurus oxyrinchus</i>	1
	Yellowtail amberjack	<i>Seriola lalandi</i>	22
October	Yellowtail amberjack	<i>Seriola lalandi</i>	3




**Figure 1.** Historical trends in catch and effort by the Canadian north Pacific albacore troll fishery from 1995 to 2011. Lines are 2-yr moving averages of catch in t (—) and effort in vessel-days (---).




**Figure 2.** Spatial distribution of the Canadian albacore troll fishery catch (number of fish) in 2011. Data are plotted on a 1° x 1° grid with symbols located on the bottom-right corner of each cell. The boundaries of the Canadian and United States exclusive economic zones (200-mile limit) are shown.



**Figure 3.** Spatial distribution of fishing effort (vessel-days) in the Canadian albacore troll fishery in 2011. Data are plotted on a 1° x 1° grid, where grid location is specified by the bottom-right corner of each grid cell. Only cells in which at least 3 vessels reported effort are shown. The black line marks the boundaries of the Canadian and United States exclusive economic zones (EEZ - 200-mile limits).




**Figure 4.** Average seasonal nominal catch-rate pattern from 2000-2009 (blue) for the Canadian albacore fishery and the 2011 seasonal catch-rate pattern. Catch-rate is calculated as the average of all 1x1 spatial strata in which fishing effort is reported during 10-day temporal strata (3 temporal strata per month). Note that the x-axis (time) is divided into three 10-day periods per month from April through November. See Kleiber and Perrin (1991) for calculation details.



**Figure 5.** Sea surface temperatures at which albacore tuna were caught by the Canadian troll fishery in 2010 (N = 940,341 fish).





**Figure 6.** Fork lengths of North Pacific albacore harvested by the Canadian fishery in 2011 by fishing zone. A total of 14,338 measurements were made in 2011.