



PLENARY 06

*21th Meeting of the
International Scientific Committee
for Tuna and Tuna-Like Species in the North Pacific Ocean
Held Virtually
July 12-21, 2021*

National Report of Japan (Japanese Tuna and Tuna-like Fisheries in the North Pacific Ocean)

Fisheries Resources Institute,
Japan Fisheries Research and Education Agency
2-12-4 Fukuura, Kanazawa, Yokohama, Kanagawa 236-8648, Japan

July 2021

Left Blank for Printing

SUMMARY

Japanese tuna fisheries consist of the three major fisheries (i.e., longline, purse seine, pole-and-line) and other miscellaneous fisheries like troll, drift-net, set-net fisheries. This paper described the recent trend of the Japanese tuna fisheries in the North Pacific Ocean and updated the statistics given in the previous National Report for ISC20 (Matsubara et al 2020). The total catch of tunas (excluding skipjack) caught by Japanese fisheries in the North Pacific Ocean was 203,521 metric ton (t) in 2019 and 132,305 t in 2020. The total catch of tunas (including skipjack) caught by Japanese fisheries in the North Pacific Ocean was 247,773 metric ton (t) in 2019 and 162,612 t in 2020. The total catch of swordfish and striped marlin was 6,114 t in 2019 and 6,897 t in 2020. In addition to fisheries description, a brief description was given on Japanese research activities on tuna and tuna-like species in the Pacific Ocean in 2020.

1 TRENDS IN FLEET SIZE

Tables 1-A and B show the number of Japanese tuna fishing vessels actually engaged in fishing by type of fishery and by vessel size class during 1980-2006 (Ministry of Agriculture, Forestry and Fishery, MAFF 1982-2008) and 2006-2020. The number of active vessels during 2006-2020 was estimated based on logbook data. The coastal longline vessels less than 20 Gross Register Tonnage (GRT), which are regulated operating only within Japan's EEZ, the research and training vessels of both longline and pole-and-line were not included in Table 1-B. The coastal longline vessels less than 20 Gross Register Tonnage (GRT), which have no license of tuna fishing and are regulated operating only within Japan's EEZ, were not included in Table 1-B. The research and training vessels of both longline and pole-and-line were not included in Table 1-B. The values of number of vessels in 2019 and 2020 were provisional in Table 1-B.

The total number of longline vessels showed a continuous declining trend since the early 1990s (Table 1-A). The number of longline vessels of the largest size class (> 200 GRT) was nearly constant in the period between the beginning of the 1980s and the mid-1990s. In accordance with the agreement of the FAO's international action plan on fishing capacity, the Japanese government implemented the fleet reduction program and decreased its large longline vessels by 20% in 1998. The number of longline vessels continued to decline thereafter. In 2009, the Japanese government implemented the second fleet reduction program for its fishery following the management measures adopted by WCPFC. Recent declining trend for the fleet size larger than 50 GRT was remarkable. The number of vessels of 100-199 GRT was 15 in 2020 which was 29% of that in 2006, and the number of vessels of 50-99 GRT was 11 in 2020 which was 25% of that in 2006 (Table 1-B). This large reduction was mainly derived from high price of fuel especially since 2007 and the fleet reduction programs implemented twice by the Government of Japan. As for the fleet size under 50 GRT, the number of vessels for 20-49 GRT showed a sharp decline since the late 1980s whereas the number of vessels of smallest size class (< 20 GRT) fluctuated at around 700 during 1980-2006 (Table 1-A). The number of vessels of 10-49 GRT was relatively stable ranging between 270 and 290 during 2006-2011 and then decreased to 202 in 2020 (Table 1-B).

The total number of purse seine vessels was 52 in 2006, and it was nearly 80% of that in the 1980s (Table 1-A). After 2006, the total number of purse seine vessels fluctuated ranging

between 67 and 75 until 2020, at a peak in 2012, 2013 and 2017. The purse seine vessels which are allowed to operate in the tropical waters are larger vessels (currently, 349 GRT or larger). The limitation of the number of such vessels has been 35 and has not changed since 1995.

The total number of pole-and-line vessels showed a continuous declining trend since 1980 (Tables 1-A and B). Suppose vessel size categories 20-49 GRT, 50-199 GRT, and over 200 GRT for 1980-2006 to compare with that for 2006-2020, the number of vessels for each category showed declining trend throughout the period (Table 1-A). The number of vessels both for 50-199 GRT and over 200 GRT showed declining trend throughout the period (Table 1-B). The number of vessels for 50-199 GRT was 36 in 2020 which is 43% of that in 2006. The number of vessels for over 200 GRT showed a declining trend until 2013, and then became stable until 2017, and then showed a declining trend again, was 22 in 2020, which is 73% of that in 2006.

2 CATCH AND EFFORT TRENDS OF THE MAJOR FISHERIES

2.1 Longline

Longline is classified by the type of license issued by the Government of Japan, i.e., coastal (< 20 GRT and can fish only in Japanese EEZ), small offshore (10-20 GRT), offshore (10-120 GRT), and distant water (> 120 GRT).

Annual distributions of fishing effort of longline in 2019 and 2020 are shown in Fig. 1. In those years, the fishing grounds were located in the east-west direction off Japan to Hawaii, the equatorial area between 15°S and 15°N, off Australia and off Peru. The fishing effort of the distant water and offshore longline remained stable at around 200 million hooks in the North Pacific in the 1980s, and then it decreased continuously to 100 million hooks in the early 2000s, and it had further decreased until 2009 (Fig. 2). After 2009, the amount of effort showed a trend of gradual decrease at a level of 35-50 million hooks.

Total catch of four tuna and four billfish species caught by distant water and offshore longline in the North Pacific has been decreased since the highest catch of 119,752 t in 1980 and was 10,508 t in 2020 which is 9% of that in 1980 (Fig. 2). Bigeye has been the dominant species in this fishery in the North Pacific. The bigeye catch, which was stable in the 1980s and about 50,000 t in the late 1980s, showed a declining trend since the 1990s, was less than 10,000 t since 2009, and was less than 5,000 since 2016. Yellowfin tuna catch ranged between 30,000 t and 50,000 t until the early 1980s. It had gradually decreased to less than 5,000 t in 2007. Albacore catch which has fluctuated around 10,000 t until 2001 decreased to about 2,000-6,000 t and kept stable at a low level during the period 2003-2020.

2.2 Purse seine

There are two types of Japanese purse seiners targeting tunas, i.e., single and group purse seine. Other than those, coastal purse seiner takes relatively small amount of tunas as a by-catch. Historically, typical group seiner consists of one purse seiner and one searching vessel and two carrier vessels, but the group seiner tends to reduce the number of vessels within each group to reduce a cost in recent years.

Fishing grounds of the purse seine were widely spreads ranging from 40°N and 10°S, from 120°E to 180°. The fishing grounds of north and south were separated by the zone from

15°N and 25°N (Fig. 3). The group seiner operates mainly in the temperate northwestern Pacific. The carrier holds fish in chilled water with ice and unloads those catches. Meanwhile, the single purse seiner (> 349 GRT) operates mainly in the tropical waters of the central and western Pacific, but a part of the vessels seasonally operates in the temperate waters.

The fishing effort and catch for the purse seine, excluding the coastal purse seine, in the North Pacific is shown in Fig. 4. The fishing effort was around 9,000 sets in the late 1980s, and then decreased to about 6,000 sets in 1998. The fishing effort generally stayed at the level about 4,000-6,000 sets in the last decade. The fishing effort was suddenly decreased to 3,320 in 2020, which is the lowest since 1980. The skipjack catch was dominant among species in this fishery, followed by yellowfin. The skipjack catch was about 150,000 t until 2008, and then decreased to 80,000 t in 2011. After 2011, the skipjack catch showed no clear trend between 80,000 t and 140,000 t. It may be exceptional in such stable situation after 2011 that the skipjack catch was 55,183 t in 2020, which correspond to the lowest level in the early 1980s.

2.3 Pole-and-line

The pole-and-line is composed of three distinct categories, i.e., coastal (< 20 GRT), offshore (10-120 GRT) and distant water (> 120 GRT) vessels in terms of the license of this fishery. Note that some of 19 GRT type vessels obtained offshore licenses since 2007, which are included in offshore category in this document. The pole-and-line can be categorized into large, middle, and small sized vessels which correspond to larger than 300 GRT, 20-300 GRT and less than 20 GRT in vessel size.

Fishing grounds of the pole-and-line were widely spread ranging from 45°N and 10°S, from 120°E to 180°. The fishing ground was rather sequential from north to south and was unlike that in the purse seine fishery. (Figs. 5). The middle-sized vessels generally operate in near shore waters of Japan and their trips are within 10 days. Southernmost fishing area for these vessels, in recent years, is near 15°N, but the important fishing ground is waters north of 25°N, around Japan and adjacent areas (Fig. 5). These vessels primarily fish skipjack and albacore from spring through autumn off the Pacific side of Japan, and harvest relatively small amount of yellowfin tuna and bigeye. They hold fish in cooled water and unload it as fresh fish. The activity of the small pole-and-line vessels is like that of the middle vessels but the area of fishing is limited within the Japanese EEZ, and the trip of these vessels is shorter. On the contrary, the large vessels tend to operate farther off waters from Japan and their trips last for two to three months. Usually they primarily target for albacore from summer through autumn season in the waters north of 20°N, and skipjack in winter and spring in the waters south of 20°N (Fig. 5). These vessels equip a brine freezer, in which fish caught are immediately stored into a tank filled with cooled brine, and then unloads it as frozen fish.

Generally, fishing effort expressed by fishing days for offshore and distant water pole-and-line rapidly decreased from around 62,000 days in the early 1980s to around 20,000 days in 1991, increased to around 23,000 days in 2000, and then gradually decreased to 7,781 days in 2020 (Fig. 6). Total catch of five tuna species for those fisheries rapidly decreased from around 280,000 t to around 170,000 t during the 1980s, and then gradually decreased from around 130,000 t to 80,000 t until the latest year (Fig. 6). Skipjack is a dominant species for this fishery, but the proportion of skipjack tends to decrease, from 80-78% in 1980-1986 to 78-60% in 2010-2020.

3 RECENT TRENDS FOR MAJOR SPECIES

3.1 Pacific bluefin tuna

Preliminary total catch of Pacific bluefin tuna (PBF) in 2020 was 7,873 t (Table 2-A), which are similar to the catch in 2019 (7,500 t). This was around the average level of past five years (2015-2019: 7,491 t). The annual catches of PBF by major fisheries in 2020 as follows; purse seine: 3,960 t, troll: 760 t, set-net: 1,234 t, longline: 1,416 t, and other fisheries: 502 t. Because of strict catch upper limit for PBF implemented since January 2015 in accordance with the WCPFC conservation and Management Measure, the annual catch since 2015 have been lower than those of the years before.

Although the most of fisheries have reduced or maintained their PBF catch since 2015, longline fishery has increased their PBF catch during the same period under the current domestic management framework. This would not be either the failures in the fishing effort control or catch control, but the increase in the availability for this fleet (Tsukahara et al., 2021). Japanese longline fishery usually catches large adult PBF, which are about 200 cm Fork Length (FL), but the length composition in recent years showed multimodal distribution which has several peaks between 110 and 200 cm FLs (Fig. 7). The highest peaks at 110 and 140 cm FLs found in 2019 and 2020 length compositions, respectively, are corresponding to the same recruitment cohort, namely 2016 year-class, which was assessed as the moderately high recruitment cohort through the stock assessment. It is also worth to note that the peaks at 110 and 160 cm FLs in 2020 composition are corresponding to the recruitments occurred between 2015 and 2017, which were fully protected cohorts by the current management measures. Increase in PBF catch by longline and its size composition possibly indicate an influx of new migrants that are smaller in the observed size as the population rebuilds.

3.2 Albacore

Preliminary total catch of albacore in 2020 was 22,249 t, which was about same as the catch in 2019 (22,243 t) and was smaller than the average of past five years (2015-2019: 35,087t). Albacore catch by the pole-and-line fluctuated largely but catch by longline did not show such fluctuations. Japanese pole-and-line fisheries target on both skipjack and albacore but particularly, large class pole-and-line vessels (> 200 GRT) primary target on albacore during summer. Recent 10 year's catch by this fishery fluctuated largely between 4,700 t and 23,000 t. Preliminary albacore catch by longline in 2020 (11,998 t) was about same as the catch in 2019 (12,216 t). Size of albacore caught by the longline fisheries in 2020 was found from 60 cm to 120 cm FL, whereas much smaller individuals (FL: 50 to 90 cm) were caught by pole-and-line (Fig. 8). No significant size differences between 2019 and 2020 were found in both fisheries.

3.3 Swordfish

Total swordfish catch in 2020 was 5,536 t which is 117% of the catches in 2019 (4,753 t). These statistics are preliminary but were smaller than the average of the past five years (2015-2019: 5,729 t). Swordfish has been caught mainly by offshore and distant water longline, which catch in 2020 was 3,266 t. The coastal longline catches in 2020 were 1,471. The Tohoku area operates night-shallow sets targeting swordfish. Longline fisheries have sampled the length composition data and are mainly distributed from 90-220 cm eye-fork length in 2019 and 2020 (Fig. 9).

3.4 Striped marlin

The total catches of striped marlin in 2020 were 1,637 t which is close to in 2019 (1,638 t). The current catches were larger than the average of the past five years (2015-2019: 1,501 t). Since the mid 1980s, striped marlin catch has been continuously decreasing primarily due to the decline of offshore and distant water longlines (from 6,378 t in 1980 to 268 t in 2020). In recent years, Japanese pelagic fisheries catch striped marlin as bycatch except for coastal drift-net and part of another longline that targets striped marlin seasonally. The length composition data of striped marlin sampled in 2019 and 2020 were distributed in the range of 100-180 cm eye-fork length (Fig. 10).

3.5 Blue shark

ISC SHARK WG conducted future projection of blue shark in 2020 after the recent three-year catches for 2016-2018 were updated using available Japanese species-specific data and statistics. Thereafter, the catch in 2019 was updated in 2021 and a total catch of blue shark in 2019 was estimated at 6,874 t. The sharp decline in 2019 was caused by decrease of catches for all fleets, especially for Offshore and Distant water longline fishery. Since there was no stock assessment of blue shark, those in 2019 were not estimated from the standardized CPUE and its fishing effort. Instead, the retained catch was used as a preliminary. Overall, those had a slightly decreasing trend since 2016 due to continuous decline of the fishing effort for longline fisheries.

3.6 Shortfin mako

ISC SHARK WG had conducted an indicator-based analysis of shortfin mako shark in 2021 and the whole-year catches for 2014-2019 of shortfin mako were updated using available Japanese species-specific data and statistics.

A total catch of shortfin mako in 2019 was estimated at 988 t. Those are stabilizing around 955-1,090 t since 2017 after decreasing those from 1,448 t in 2016. This is due to the decline of catch for Offshore and Distant water longline fishery as well as for large mesh drift-net fishery.

3.7 Others (Bigeye, Skipjack and Yellowfin tunas)

Preliminary total catch of bigeye in 2020 was 14,567 t which corresponds to 4% increase of the catch in 2019 (14,055 t) and was lower than the average of past five years (2015-2019: 16,015 t). Total catch of bigeye by Japanese fisheries showed no clear trend in the last six years and longline has been the highest proportion among gears in the North Pacific.

Preliminary total catch of skipjack in 2020 was 87,671 t which corresponds to 45% decrease of the catch in 2019 (159,523 t) and was lower than the average of past five years (2015-2019: 173,267 t). Most of skipjack was caught by pole-and-line and purse seine in the North Pacific. Total catch of skipjack by Japanese fisheries showed no clear trend in the last six years in the North Pacific. The lower total catch in 2020 were due to lower catch by both pole-and-line and purse seine fisheries.

Preliminary total catch of yellowfin tuna in 2020 was 30,307 t which corresponds to 32% decrease of the catch in 2019 (44,213 t) and was lower than the average of past five years

(2015-2019: 44,252 t). The yellowfin tuna caught by purse seine has been the highest proportion among gears in the North Pacific. The lower total catch in 2020 were due to lower catch by the purse seine fishery in the tropical waters in the North Pacific).

4 RESEARCH ACTIVITIES

The Fisheries Resources Institute (FRI) of Japan and local prefectural fisheries experimental stations, has run the nationwide port sampling project for collection of catch, effort and size data at the major landing ports since the early 1990s. The tagging studies using conventional, archival and popup satellite archival tags have been conducted by research and training vessels as well as commercial vessels. In addition, there were cooperative works with prefectural fisheries experimental stations and universities. Several cooperative studies were also conducted with foreign countries.

4.1 Pacific Bluefin tuna

4.1.1 *Pacific bluefin tuna larvae/juveniles research cruise*

Since 2011, larval and juvenile surveys have been conducted to estimate current main spawning area and period of PBF. In 2020, research cruises were designed to focus on ecological studies of larval/juvenile PBF by R/Vs Shunyo-Maru, Yoko-Maru, Hokko-Maru and five prefectural R/Vs. Larval surveys were conducted in the south of Japan around Nansei Islands area, where is a major spawning ground of PBF, from May to August and also in the Sea of Japan, which is another spawning ground of PBF, from July to August. In addition to these two spawning grounds, larval survey was conducted in Joban area in the coastal area of northeastern Japan in July and August. In 2020, over 650 of PBF larvae were captured in the spawning grounds. Small juveniles of PBF around 2-5 cm FL were also captured in Nansei Island area and Joban area by small surface-trawl net. Juvenile surveys were conducted in nursery areas in the Sea of Japan in September. Over 430 of PBF juveniles were captured by surface-trawl net in the Sea of Japan in 2020.

Collected samples are being examined by a variety of approaches such as genetic identification, aging, growth analysis, stable isotope, microchemistry and stomach contents analyses to elucidate the survival processes of larval and juvenile PBF in relation to biological and environmental factors, which should help to understand the recruitment mechanism to PBF fisheries around Japan.

4.1.2 *Troll survey on age-0 Pacific Bluefin tuna*

Recruitment abundance index (standardized CPUE of age-0 PBF) for current stock assessment is based on the sales slips of Japanese troll fishery in the East China Sea. The FRI has also conducted monitoring survey of troll fishery's operations to obtain alternative indices, which are targeting age-0 PBF. Currently, the data logger and transmitter were installed on 77 troll fishing boats in six prefectures (Mie, Wakayama, Kochi, Miyazaki, Nagasaki and Shimane). Once the fishermen input their catch number of age-0 fish during the fishing operation, the catch information with geographic position and Sea Surface Temperature (SST) data are sent to the FRI via a cellular network on a real time basis.

The data from 67 boats in 2020 were available for evaluation of recruitment. Although the data coverage over the entire troll operations by this survey are lower than that of the conventional troll CPUE which is used for the current assessment, those operational and environmental information in finer spatial-temporal resolutions enable to analyze the recruitment trend of PBF in details. The FRI reported the preliminary recruitment indices, which was standardized CPUE obtained through this monitoring system, to the PBFWG meeting of the ISC (Nishikawa et al., 2021). In the PBFWG meeting held in April 2021, the feasibility of this index as an input data for the assessment was discussed, and the WG agreed to continue their investigation on this index for the future stock assessment.

4.1.3 Size sampling for Pacific Bluefin tuna farming operation

Since 2017, stereoscopic cameras have been introduced in the bluefin ranch sector in Japan and are utilized to obtain counts and estimates of individual length and weight of penned PBF by the Japanese Small Pelagic Fish Purse Seine fleet (JSPF-PS). The FRI conducted an experiment to validate (or not) the reliability of the length data obtained by the stereoscopic cameras and concluded that the length composition data obtained by this new technology can provide a large volume of high-quality length data (Fukuda and Nakatsuka, 2019). Those data were included in the latest stock assessment to depict the number and size of PBF removed by the JSPF-PS operated for the farming. In 2020, all the penning operations (83 operations in total) by the JSPF-PS were measured using this technology, and consequently, 3.9 % of the total penned PBF were measured.

4.1.4 Tissue sampling and technical development for close-kin analysis

Tissue sampling for close-kin analysis has started since 2015 and around 2,000 individuals were sampled in 2020. Large mature adults of PBF (about 120-300 kg in BW (Body Weight)) were sampled from individuals fished by coastal longline fishery around Okinawa Islands in late April to early July, while young-of-the-year juveniles (about 0.1-0.3 kg in BW) caught by troll fishery during summer in the Pacific coastal water off Western Japan, such as Kochi Prefecture. The breeding or hatching area of these samples can be identified as water around Nansei archipelago based on the fishing timing and area. For the other spawning ground, The Sea of Japan, young adults and either nearly-matured or matured individuals (about 20-60 kg in BW) were sampled from fish landed at Sakai-minato, while young-of-the-year juveniles were sampled in Oki islands. Additionally, there are samples a lot of samples to assure the randomness of sampling from all over Japan, especially in the Tsugaru straits. All of muscle tissues sampled were preserved in specific buffer (TNES-Urea 6M buffer) because of the higher stability of content DNA.

The close-kin project team, which consists of FRI and collaborating academic researchers, has been working to develop practical procedures of close-kin analysis in PBF. The team began to conduct genotyping of actual samples since 2018 using random amplicon sequencing method. Alongside, the team is addressing development of statistical approaches for estimating stock abundance. Furthermore, a numerical model which can keep all genealogical information among every individual has been developed with expecting application to the operating model of statistical approach and enhancement in the quality of sampling design.

4.2 Sharks, Swordfish, and billfishes

4.2.1 Port sampling and the onboard research program for sharks in Kesenuma fishing port

In 2020 size and sex data of both blue shark and shortfin mako shark were collected from port sampling in Kesenuma fishing port, located in northeastern Honshu (the main island of Japan), and the onboard research project for Kesenuma offshore longline fleet throughout the year. In the port sampling, size and data from 36,093 blue shark were collected and 71% of individuals measured were males. In addition, 68% of males and 65% of females were juveniles.

Regarding blue shark, the number of catch by four size categories was recorded in the onboard research program for Kesenuma-offshore longline fleet. Total of 160,097 blue sharks were recorded by size category and “large (processed weight: ≥ 15 kg)” consisted 52% of all catch with 27% of “middle ($11 \text{ kg} \leq \text{processed weight} < 15 \text{ kg}$)”, 19% of “small ($5 \text{ kg} \leq \text{processed weight} < 11 \text{ kg}$)” and 2% of “extra small (processed weight: $< 5 \text{ kg}$)”.

For shortfin mako, size data from 7,050 individuals were collected from port sampling (male: 3,069, female:2,955, unknown:1,026) and 64% of males and 100% of females were juveniles. In contrast to blue shark, almost all of sampled female were juvenile in shortfin mako. Total of 8,481 sharks was recorded by size category from the onboard research by Kesenuma-offshore longline fleet and “large (precaudal length: $> 200 \text{ cm}$)” consisted 5% of all catch with 34% of “middle ($150 \text{ cm} < \text{precaudal length} \leq 200 \text{ cm}$)”, 47% of “small ($100 \text{ cm} < \text{precaudal length} \leq 150 \text{ cm}$)” and 13% of “extra small (precaudal length $\leq 100 \text{ cm}$)”.

4.2.2 Tagging study for sharks

In the research cruise by No.37 Den-Maru (chartered longline vessel) conducted between 17th April and 28th May in 2020, normal tag was attached to 235 blue sharks and 21 shortfin mako. Two blue sharks were recaptured 33 and 49 days after release.

In the research cruise research and training vessel (RTV), normal tag was attached to 23 blue sharks.

In 2020, FRI and NOAA (Pacific Islands Fisheries Science Center) launched the collaborative study on the migration of blue shark in the central North Pacific using popup satellite archival tag (PSAT). Due to the influence of COVID-19, deployment of PSAT has been still ongoing.

4.2.3 Biological sample collection for sharks, swordfish and billfishes

Whole body of 32 swordfish and 9 striped marlin were collected by chartered longline vessel (No. 37 Den-Maru and No. 188 Han ei-Maru), RTV and commercial longline vessel. In addition, head, internal organ including testes/ovary, and the first dorsal/anal fin were collected from 6 swordfish and 2 striped marlins during the research cruise by No. 37 Den-Maru for the study of their life history parameters.

For the analysis of genetic population structure of billfish, the whole body of juvenile swordfish (N=17) and striped marlin (N=1) were collected from Kesenuma offshore longline fleet.

For the improvement life history parameter and ecological study of sharks, muscle tissue, reproductive organ, stomach were collected from 3 blue shark, 8 shortfin mako (whole body for 3 mako). Blood samples were collected from 8 blue sharks for reproductive/genetic analysis.

All of the samples above were collected by the research cruise and commercial/training longline operation conducted in the North Pacific in 2020.

4.3 Skipjack

4.3.1 *Tagging for Skipjack*

The FRI has been conducting skipjack tagging research mainly to investigate migration patterns to the fishing ground off Japan. One offshore pole-and-line vessel (20-119 GRT) and one distant water pole-and-line vessel (> 199 GRT) were fully chartered to conduct the research off Japan in October 2020 and in tropical areas (5°-25°N, 140°-180°E) in December 2020, respectively. A total of 10,074 skipjack tuna (2,774 off Japan and 7,300 in tropical areas) were released including 491 individuals (160 off Japan and 331 in tropical areas) with archival tags (Lotek LAT2910). In addition, skipjack tagging has been conducted in cooperation with Ajinomoto Co., Inc. in the coastal area of southwestern Japan since 2009. In 2020, skipjack tuna tagging was planned, however, the tagging was postponed due to the COVID-19 pandemic.

Besides above studies, five research/training cruises on pole-and-line vessels conducted skipjack tagging in 2020 around Japanese water. A total of 500 skipjack tuna including 116 individuals with archival tags were released in the south off Japan, around Izu Islands, around Hachijo Island (33°N, 139°E), and Wakayama (33.15°N, 135.75°E).

4.4 Albacore

4.4.1 *Tagging for Albacore*

The FRI started new tagging research to investigate female and male albacore distribution and migration in the north western Pacific. In the 2020 research, 19 and 8 archival tags were attached to albacore in off Kushimoto, Wakayama in February 2020, and off Kume islands, Okinawa in November 2020, respectively.

5 REFERENCE

- Ohashi, Y., Tsukahara, Y., and Nakatsuka, S. (2019) Japanese coastal longline CPUE for Pacific bluefin tuna: Preliminary update up to 2018 fishing year: ISC/19/PBFWG-2/09.
- Fukuda, H. and Nakatsuka, S. (2019) Estimation of the PBF length-composition for the Japanese purse seine with new data collected at young PBF farming operation using stereoscopic camera: ISC/19/PBFWG-2/02.
- MAFF (1982-2008): Gyogyou yousyokugyou seisan toukei nenpou (Yearbook of fisheries and aquaculture production statistics of Japan for 1980-2006), Statistics Department, Minister's Secretariat, Ministry of Agriculture, Forestry and Fishery.
- Tsukahara, Y., Okamoto, H., Chiba, K., Ohashi, Y., Kiyofuji, H., Ijima H., Kai, M., Semba, Y., Aoki, Y., Fujioka, K., Tawa, A., Fukuda, H., Ohashi, S., and Uosaki, K. (2019) National report of Japan. ISC/19/Plenary/06. 22pp.

6 TABLE

Table 1. Number of Japanese tuna fishing vessels.

A. Number of Japanese tuna fishing vessels operated in the Pacific Ocean by type of fisheries and vessel size based on MAFF (1980-2006).

| Year | Longline fishery ^{*1} | | | | | Total | Purse seine fishery | | | Pole-and-line fishery | | | | | Total |
|------|--------------------------------|-----------|-----------|-------------|---------|-------|--------------------------|---------|-------|------------------------|-----------|-----------|-------------|---------|-------|
| | 1-19 GRT | 20-49 GRT | 50-99 GRT | 100-199 GRT | 200-GRT | | 50-199 GRT ^{*2} | 200-GRT | Total | 1-19 GRT ^{*3} | 20-49 GRT | 50-99 GRT | 100-199 GRT | 200-GRT | |
| 1980 | 821 | 57 | 715 | 103 | 645 | 2,341 | 50 | 16 | 66 | 3,232 | 14 | 350 | 10 | 198 | 3,804 |
| 1981 | 774 | 55 | 706 | 100 | 661 | 2,296 | 50 | 23 | 73 | 3,064 | 10 | 353 | 6 | 179 | 3,612 |
| 1982 | 722 | 43 | 634 | 90 | 589 | 2,078 | 52 | 33 | 85 | 3,011 | 11 | 320 | 6 | 138 | 3,486 |
| 1983 | 561 | 38 | 589 | 93 | 550 | 1,831 | 59 | 36 | 95 | 3,021 | 12 | 297 | 9 | 116 | 3,455 |
| 1984 | 523 | 32 | 538 | 108 | 610 | 1,811 | 54 | 33 | 87 | 2,904 | 8 | 273 | 10 | 105 | 3,300 |
| 1985 | 620 | 28 | 512 | 131 | 628 | 1,919 | 47 | 35 | 82 | 2,754 | 8 | 244 | 9 | 95 | 3,110 |
| 1986 | 536 | 25 | 435 | 168 | 632 | 1,796 | 53 | 38 | 91 | 2,455 | 6 | 224 | 9 | 91 | 2,785 |
| 1987 | 661 | 23 | 348 | 197 | 649 | 1,878 | 47 | 34 | 81 | 2,404 | 6 | 210 | 9 | 89 | 2,718 |
| 1988 | 586 | 21 | 289 | 233 | 649 | 1,778 | 48 | 39 | 87 | 2,613 | 5 | 191 | 11 | 70 | 2,890 |
| 1989 | 650 | 20 | 248 | 238 | 653 | 1,809 | 43 | 37 | 80 | 2,254 | 3 | 187 | 12 | 67 | 2,523 |
| 1990 | 685 | 21 | 227 | 241 | 664 | 1,838 | 43 | 35 | 78 | 2,228 | 4 | 176 | 9 | 66 | 2,483 |
| 1991 | 768 | 19 | 199 | 222 | 682 | 1,890 | 38 | 35 | 73 | 2,277 | 3 | 166 | 10 | 63 | 2,519 |
| 1992 | 793 | 19 | 164 | 206 | 681 | 1,863 | 31 | 38 | 69 | 2,093 | 3 | 156 | 11 | 46 | 2,309 |
| 1993 | 790 | 18 | 138 | 201 | 682 | 1,829 | 27 | 36 | 63 | 1,927 | 3 | 147 | 10 | 43 | 2,130 |
| 1994 | 819 | 21 | 110 | 198 | 675 | 1,823 | 23 | 33 | 56 | 1,830 | 3 | 124 | 10 | 48 | 2,015 |
| 1995 | 738 | 20 | 92 | 187 | 667 | 1,704 | 20 | 31 | 51 | 481 | 3 | 104 | 20 | 46 | 654 |
| 1996 | 711 | 17 | 91 | 155 | 640 | 1,614 | 21 | 32 | 53 | 512 | 3 | 89 | 29 | 43 | 676 |
| 1997 | 698 | 11 | 88 | 145 | 631 | 1,573 | 20 | 35 | 55 | 436 | 2 | 76 | 39 | 45 | 598 |
| 1998 | 712 | 11 | 80 | 129 | 623 | 1,555 | 20 | 35 | 55 | 382 | 2 | 73 | 40 | 46 | 543 |
| 1999 | 703 | 6 | 78 | 119 | 567 | 1,473 | 22 | 36 | 58 | 416 | 1 | 62 | 54 | 46 | 579 |
| 2000 | 732 | 3 | 76 | 111 | 496 | 1,418 | 23 | 37 | 60 | 357 | 1 | 56 | 57 | 47 | 518 |
| 2001 | 777 | 4 | 76 | 110 | 494 | 1,461 | 19 | 36 | 55 | 285 | 1 | 49 | 59 | 47 | 441 |
| 2002 | 780 | 4 | 69 | 110 | 484 | 1,447 | 18 | 36 | 54 | 251 | 1 | 45 | 58 | 48 | 403 |
| 2003 | 764 | 3 | 64 | 99 | 460 | 1,390 | 17 | 36 | 53 | 292 | 1 | 44 | 56 | 44 | 437 |
| 2004 | 702 | 2 | 55 | 77 | 455 | 1,291 | 17 | 36 | 53 | 284 | 1 | 38 | 57 | 43 | 423 |
| 2005 | 694 | 2 | 46 | 59 | 432 | 1,233 | 17 | 36 | 53 | 247 | 1 | 36 | 58 | 45 | 387 |
| 2006 | 709 | 1 | 43 | 54 | 401 | 1,208 | 16 | 36 | 52 | 213 | 1 | 27 | 58 | 36 | 335 |

^{*1} Longline vessels larger than 50 GRT include those operated in the area other than the Pacific^{*2} 50-199 GRT class vessels only include those operated in the Pacific side of northern Japan.^{*3} 1-19 GRT class vessels before 1995 include those engaged in trolling

B. Number of Japanese tuna fishing vessels operated in the North Pacific Ocean by type of fisheries and vessel size based on logbook. Values in 2019 and 2020 are provisional.

| Year | Longline fishery | | | | | Purse seine fishery | | | | Pole-and-line fishery | | | |
|------|------------------|--------------|----------------|-------------|-------|----------------------------|----------------|-------------|-------|-----------------------|---------------|-------------|-------|
| | 10-49 GRT | 50-99 GRT | 100-199 GRT | 200- GRT | Total | 50-199 GRT ² | 200-499 GRT | 500- GRT | Total | 20-49 GRT | 50-199 GRT | 200- GRT | Total |
| 2006 | 277 | 44 | 52 | 113 | 486 | 31 | 35 | 1 | 67 | 1 | 83 | 30 | 114 |
| 2007 | 279 | 42 | 48 | 89 | 458 | 34 | 36 | 1 | 71 | 1 | 77 | 29 | 107 |
| 2008 | 277 | 42 | 40 | 90 | 449 | 35 | 37 | 1 | 73 | 1 | 69 | 29 | 99 |
| 2009 | 277 | 38 | 33 | 81 | 429 | 33 | 36 | 3 | 72 | 1 | 68 | 28 | 97 |
| 2010 | 290 | 29 | 28 | 98 | 445 | 31 | 35 | 4 | 70 | 1 | 66 | 28 | 95 |
| 2011 | 273 | 24 | 25 | 99 | 421 | 33 | 36 | 4 | 73 | 0 | 63 | 28 | 91 |
| 2012 | 265 | 21 | 21 | 92 | 399 | 34 | 37 | 4 | 75 | 0 | 60 | 27 | 87 |
| 2013 | 260 | 20 | 23 | 87 | 390 | 34 | 37 | 4 | 75 | 0 | 55 | 25 | 80 |
| 2014 | 250 | 18 | 21 | 90 | 379 | 33 | 37 | 3 | 73 | 1 | 54 | 25 | 80 |
| 2015 | 239 | 18 | 24 | 80 | 361 | 30 | 35 | 5 | 70 | 1 | 51 | 24 | 76 |
| 2016 | 234 | 16 | 16 | 64 | 330 | 32 | 33 | 4 | 69 | 1 | 50 | 25 | 76 |
| 2017 | 233 | 15 | 15 | 50 | 313 | 37 | 34 | 4 | 75 | 1 | 48 | 31 | 80 |
| 2018 | 229 | 14 | 16 | 52 | 311 | 34 | 30 | 4 | 68 | 1 | 44 | 25 | 70 |
| 2019 | 230 | 13 | 17 | 44 | 304 | 36 | 31 | 5 | 72 | 1 | 42 | 24 | 67 |
| 2020 | 202 | 11 | 15 | 43 | 271 | 33 | 30 | 6 | 69 | 1 | 36 | 22 | 59 |

*4 50-199 GRT class vessels only include those operated in the Pacific side of northern Japan.

Table 2. Catch in weight (t) by species by fisheries in the North Pacific.**A. Pacific bluefin tuna**

| Year | Purse Seine | | Dist. & Off. Longline ^{*5} | | Coastal Longline ^{*5} | Troll | Pole and Line | Set Net | Others ^{*7} | Total |
|--------------------|-------------|----------|-------------------------------------|---------------|--------------------------------|-------|---------------|---------|----------------------|-------|
| | Tuna PS | Small PS | North Pacific | South Pacific | | | | | | |
| 2015 | 2,759 | 886 | 11 | 4 | 637 | 413 | 8 | 1,242 | 431 | 6,390 |
| 2016 | 3,267 | 1,828 | 14 | 4 | 677 | 778 | 54 | 1,228 | 508 | 8,359 |
| 2017 | 3,341 | 1,199 | 21 | 6 | 892 | 605 | 49 | 2,221 | 665 | 9,000 |
| 2018 | 3,225 | 825 | 21 | 0 | 679 | 371 | 9 | 645 | 431 | 6,205 |
| 2019 | 3,213 | 1,251 | 25 | 0 | 977 | 720 | 0 | 941 | 372 | 7,500 |
| 2020 ^{*6} | 3,208 | 752 | 75 | 0 | 1,341 | 760 | 1 | 1,234 | 502 | 7,873 |

*5: Distant-water and Offshore longline vessels are mainly 20 GRT or larger, and most of coastal longline vessels are smaller than 20 GRT.

*6: Most recent year's catch value is provisional.

*7: Others include drift net, handline, trawl, other longline, and unclassified fisheries.

B. Albacore

| Year | Longline | | Pole-and-line | | | Drift-net | Purse seine | Troll | Set-net | Others | Total |
|------|--|----------|----------------|----------|---------|-----------|-------------|-------|---------|--------|----------|
| | Distant Water + Offshore ^{*8} | Coastal | Distant Waters | Offshore | Coastal | | | | | | |
| 2015 | 3,907 | 17,106 | 11,498 | 9,710 | 86 | 138 | 1,072 | 239 | 17 | 167 | 43,940 |
| 2016 | 3,431 | 13,118 | 8,648 | 5,754 | 33 | 19 | 3,679 | 148 | 28 | 128 | 34,986 |
| 2017 | 3,710 | 13,598 | 12,108 | 8,753 | 30 | 40 | 1,250 | 107 | 48 | 119 | 39,763 |
| 2018 | 3,070 | 10,121 | 9,362 | 8,394 | 119 | 35 | 3,039 | 78 | 13 | 70 | 34,301 |
| 2019 | 2,906 | 9,310 | 4,669 | 3,662 | 177 | 9 | 1,045 | 543 | 27 | 95 | 22,443 |
| 2020 | (1,658) | (10,341) | (4,693) | (3,662) | (177) | (9) | (1,045) | (543) | (27) | (95) | (22,249) |

*8 Category distant water + Offshore longline vessels include training/research vessels.

() Different data source or carry over from previous year.

C. Swordfish

| Year | Longline | | | | | Drift-net | Bait fishing | Net fishing | Trap-net | Others | Total |
|--------------------|--------------------------|---------|--------|-----------|--------------|-----------|--------------|-------------|----------|--------|-------|
| | Distant Water + Offshore | Coastal | Others | Drift-net | Bait fishing | | | | | | |
| 2015 | 3,755 | 1,343 | 1 | 277 | 281 | 0 | 3 | 204 | 5,865 | | |
| 2016 | 3,509 | 2,094 | 2 | 303 | 256 | 0 | 2 | 169 | 6,335 | | |
| 2017 | 2,860 | 1,975 | 2 | 291 | 289 | 0 | 3 | 274 | 5,694 | | |
| 2018 | 3,212 | 1,801 | 2 | 230 | 267 | 0 | 5 | 480 | 5,997 | | |
| 2019 ^{*9} | 2,611 | 1,343 | 2 | 242 | 210 | 0 | 6 | 339 | 4,753 | | |
| 2020 ^{*9} | 3,266 | 1,471 | 2 | 242 | 210 | 0 | 6 | 339 | 5,536 | | |

*9 Catch between 2019 and 2020 are preliminary.

D. Striped Marlin

| Year | Longline | | | Drift-net | Bait fishing | Net fishing | Trap-net | Others | Total |
|---------------------|--------------------------|---------|--------|-----------|--------------|-------------|----------|--------|-------|
| | Distant Water + Offshore | Coastal | Others | | | | | | |
| 2015 | 267 | 1,131 | 43 | 287 | 27 | - | 37 | 37 | 1,829 |
| 2016 | 245 | 778 | 33 | 308 | 32 | 0 | 25 | 41 | 1,462 |
| 2017 | 160 | 764 | 53 | 241 | 28 | - | 28 | 23 | 1,296 |
| 2018 | 147 | 711 | 28 | 278 | 36 | - | 28 | 52 | 1,280 |
| 2019 ^{*10} | 255 | 984 | 29 | 241 | 39 | - | 29 | 61 | 1,638 |
| 2020 ^{*10} | 268 | 970 | 29 | 241 | 39 | - | 29 | 61 | 1,637 |

*10 Catch between 2019 and 2020 are preliminary.

E. Blue shark

| Year | Longline | | | | Large mesh drift-net | Bait fishing | Trap-net | Others | Total |
|---------------------|---------------|----------|---------|--------|----------------------|--------------|----------|--------|--------|
| | Distant Water | Offshore | Coastal | Others | | | | | |
| 2014 | 6,392 | 8,714 | 538 | 598 | 1,060 | 2 | 4 | 0 | 17,306 |
| 2015 | 6,163 | 5,370 | 551 | 386 | 1,080 | 2 | 21 | 0 | 13,573 |
| 2016 | 8,874 | 5,008 | 375 | 225 | 1,832 | 2 | 26 | 1 | 16,343 |
| 2017 | 8,611 | 5,621 | 342 | 212 | 1,366 | 1 | 4 | 0 | 16,158 |
| 2018 | 5,712 | 4,787 | 263 | 159 | 1,236 | 1 | 40 | 0 | 12,198 |
| 2019 ^{*11} | 2,558 | 2,862 | 208 | 165 | 1,052 | 1 | 28 | 0 | 6,874 |

*11 Values of Distant water longline and Offshore longline are retained catches and preliminary.

F. Shortfin mako

| Year | Longline | | | Large mesh drift-net | Trap-net and others | Total |
|------|--|---------------------------------------|-------------------|----------------------|---------------------|-------|
| | Offshore and Distant water (Shallow set) | Offshore and Distant water (Deep set) | Coastal and other | | | |
| 2014 | 609 | 139 | 7 | 263 | 3 | 1,021 |
| 2015 | 605 | 240 | 2 | 334 | 11 | 1,193 |
| 2016 | 784 | 171 | 32 | 446 | 16 | 1,448 |
| 2017 | 564 | 87 | 23 | 271 | 10 | 955 |
| 2018 | 638 | 181 | 19 | 223 | 28 | 1,090 |
| 2019 | 571 | 204 | 16 | 195 | 2 | 988 |

G. Bigeye

| Year | Longline | Pole-and-line | Purse seine | Gillnet | Set-net | Troll | Other | Total |
|------|----------|---------------|-------------|---------|---------|-------|-------|--------|
| 2015 | 16,030 | 780 | 2,855 | 4 | 3 | 140 | 114 | 19,926 |
| 2016 | 11,598 | 1,012 | 1,310 | 0 | 1 | 87 | 109 | 14,117 |
| 2017 | 11,425 | 1,444 | 2,201 | 1 | 0 | 119 | 89 | 15,279 |
| 2018 | 11,631 | 1,432 | 3,471 | 1 | 0 | 80 | 84 | 16,699 |
| 2019 | 11,838 | 549 | 1,444 | 1 | 0 | 110 | 113 | 14,055 |
| 2020 | 11,763 | 959 | 1,621 | 1 | 0 | 110 | 113 | 14,567 |

H. Skipjack

| Year | Longline | Pole-and-line | Purse seine | Gillnet | Set-net | Troll | Other | Total |
|------|----------|---------------|-------------|---------|---------|-------|-------|---------|
| 2015 | 77 | 70,353 | 117,548 | 119 | 153 | 1,238 | 46 | 189,534 |
| 2016 | 34 | 68,981 | 82,658 | 111 | 264 | 1,642 | 53 | 153,743 |
| 2017 | 30 | 61,940 | 93,396 | 61 | 401 | 1,615 | 81 | 157,524 |
| 2018 | 21 | 78,998 | 125,119 | 91 | 494 | 1,154 | 133 | 206,010 |
| 2019 | 38 | 73,592 | 84,054 | 96 | 246 | 1,387 | 110 | 159,523 |
| 2020 | 23 | 30,524 | 55,285 | 96 | 246 | 1,387 | 110 | 87,671 |

I. Yellowfin tuna

| Year | Longline | Pole-and-line | Purse seine | Gillnet | Set-net | Troll | Other | Total |
|------|----------|---------------|-------------|---------|---------|-------|-------|--------|
| 2015 | 9,147 | 2,971 | 25,503 | 12 | 56 | 2,014 | 599 | 40,302 |
| 2016 | 7,978 | 3,218 | 21,300 | 16 | 120 | 2,250 | 806 | 35,688 |
| 2017 | 7,966 | 3,201 | 24,195 | 7 | 135 | 1,877 | 690 | 38,071 |
| 2018 | 7,955 | 3,519 | 38,868 | 6 | 77 | 1,738 | 587 | 52,750 |
| 2019 | 11,223 | 2,930 | 27,039 | 4 | 208 | 2,070 | 778 | 44,252 |
| 2020 | 7,212 | 2,344 | 17,691 | 4 | 208 | 2,070 | 778 | 30,307 |

7 FIGURES

Longline fishery




Fig 1. Distribution of fishing effort (Number of hooks) for the Japanese distant water and offshore longline fisheries in the Pacific, 2019-2020.




Fig 2. Historical catches in weight (t) for major species and fishing effort (Number of hooks in million) of the Japanese distant water and offshore longline fisheries (not including small offshore fishery) in the North Pacific Ocean. PBF: Pacific bluefin tuna, ALB: albacore, BET: bigeye, YFT: yellowfin tuna, SWO: swordfish, MLS:

striped marlin, BUM: blue marlin, BLM: black marlin. Values in 2019 and 2020 are provisional.

Purse seine fishery




Fig 3. Distribution of fishing effort (number of sets) for the Japanese purse seine fishery in the Pacific, 2019-2020.




Fig 4. Historical catches in weight (t) for major species and fishing effort (Number of sets) of the Japanese purse seine fishery in the North Pacific Ocean. SKJ: skipjack, YFT: yellowfin tuna, BET: bigeye, PBF: Pacific bluefin tuna, ALB: albacore. Value in 2020 is provisional.

Pole-and -line fishery




Fig 5. Distribution of fishing effort (number of days) of the Japanese pole-and-line fishery (larger than 20 GRT vessels) in the Pacific, 2019-2020.




Fig 6. Historical catches in weight (t) for major species and fishing effort (Number of fishing days) of Japanese distant water and offshore fisheries in the North Pacific. SKJ: skipjack, ALB: albacore, YFT: yellowfin tuna, PBF: Pacific bluefin tuna, BET: bigeye. The catch for PBF includes the catch by coastal pole-and-line (less than 20 GRT vessels) fishery. Value in 2020 is provisional.




Fig 7. Annual relative length frequency distribution (simply summing up all measurements) for Pacific bluefin tuna (PBF) caught by longline in 2019 (left) and 2020 (right). Texts in each graph indicate gear, species, year, and the number of fish measured.




Fig 8. Annual relative length frequency distribution (simply summing up all measurements) for albacore (ALB) caught by longline (upper two panels) and pole-

and-line (lower two panels) in 2019 (left) and 2020 (right). Texts in each graph indicate gear, species, year, and the number of fish measured.




Fig 9. Annual relative length frequency distribution (simply summing up all measurements) for swordfish (SWO) caught by longline in 2019 (left) and 2020 (right). Texts in each graph indicate gear, species, year, and the number of fish measured.




Fig 10. Annual relative length frequency distribution (simply summing up all measurements) for striped marlin (MLS) caught by longline in 2019 (left) and 2020 (right). Texts in each graph indicate gear, species, year, and the number of fish measured.

Fig 11. fish measured.