

FINAL

ISC/24/ PLENARY/06



PLENARY 06

*24th Meeting of the
International Scientific Committee for Tuna
and Tuna-Like Species in the North Pacific Ocean
Victoria, Canada
June 19-24, 2024*

NATIONAL REPORT OF JAPAN: JAPANESE TUNA AND TUNA-LIKE FISHERIES IN THE NORTH PACIFIC OCEAN IN 2023

Fisheries Resources Institute,
Japan Fisheries Research and Education Agency
2-12-4 Fukuura, Kanazawa, Yokohama, Kanagawa 236-8648, Japan

Kirara Nishikawa, Hidetada Kiyofuji, Shuya Nakatsuka, Hiromu Fukuda, Ryuichi Matsukura,
Yuichi Tsuda, Takayuki Matsumoto, Hiroataka Ijima, Marko Jusup, Yasuko Semba, Mikihiko
Kai, Yoshinori Aoki, Kei Okamoto, Naoto Matsubara, Takaaki Hasegawa, Yohei Tsukahara, and
Keisuke Satoh

June 2024

Left Blank for Printing

SUMMARY

Japanese tuna fisheries consist of the three major fisheries (i.e., longline, purse seine, and pole-and-line) and other miscellaneous fisheries like troll, driftnet, and setnet fisheries. This paper described the recent trend of the Japanese tuna fisheries in the North Pacific Ocean and updated the statistics given in the previous National Report for ISC23 (Kai et al., 2023). The total catch of tunas excluding skipjack caught by Japanese fisheries in the North Pacific Ocean was 61,235 metric ton (t) in 2022 and 65,055 t in 2023. The total catch of tunas including skipjack caught by Japanese fisheries in the North Pacific Ocean was 181,620 t in 2022 and 177,371 t in 2023. The total catch of swordfish and striped marlin was 5,287 t in 2022 and 6,515 t in 2023. In addition to fisheries description, a brief description was given on Japanese research activities in 2023 for tuna and tuna-like species in the Pacific Ocean.

1. TRENDS IN FLEET SIZE

Tables 1-A and B show the number of Japanese tuna fishing vessels engaged in fishing by type of fishery and by vessel size class during 1980-2006 (Ministry of Agriculture, Forestry and Fishery, MAFF 1982-2008) and 2006-2023. The number of active vessels during 2006-2023 was estimated based on logbook data. The coastal longline vessels less than 20 Gross Register Tonnage (GRT), which are regulated operating only within Japan's Exclusive Economic Zone (EEZ), the research and training vessels of both longline and pole-and-line were not included in Table 1-B. The coastal longline vessels less than 20 GRT, which have no license of tuna fishing and are regulated operating only within Japan's EEZ, were not included in Table 1-B. The research and training vessels of both longline and pole-and-line were not included in Table 1-B. The values of number of vessels in 2022 and 2023 were provisional in Table 1-B.

The total number of longline vessels showed a continuous declining trend since the early 1990s (Table 1-A). The number of longline vessels of the largest size class (> 200 GRT) was nearly constant in the period between the beginning of the 1980s and the mid-1990s. In accordance with the agreement of the Food and Agriculture Organization of the United Nations (FAO)'s international action plan on fishing capacity, the Japanese government implemented the fleet reduction program and decreased its large longline vessels by 20% in 1998. The number of longline vessels continued to decline thereafter. In 2009, the Japanese government implemented the second fleet reduction program for its fishery following the management measures adopted by the Western and Central Pacific Fisheries Commission (WCPFC). The recent declining trend for the fleet size larger than 50 GRT was remarkable. The number of vessels of 50-99 GRT was 6 in 2023 which is 13% of that in 2006, and the number of vessels of 100-199 GRT was 13 in 2023 which is 25% of that in 2006 (Table 1-B). This large reduction was mainly derived from high price of fuel especially since 2007 and the fleet reduction programs implemented twice by the Government of Japan. As for the fleet size under 50 GRT, the number of vessels for 20-49 GRT showed a sharp decline since the late 1980s whereas the number of vessels of smallest size class (< 20 GRT) fluctuated at around 700 during 1980-2006 (Table 1-A). The number of vessels of 10-49 GRT was relatively stable ranging between 273 and 290 during 2006-2011 and then decreased to 195 in 2023 (Table 1-B).

The total number of purse seine vessels was 52 in 2006, and it was nearly 80% of that in the 1980s (Table 1-A). After 2006, the total number of purse seine vessels fluctuated ranging between 53 and 75 until 2023. The purse seine vessels which are allowed to operate in the tropical waters are

larger vessels (currently, 349 GRT or larger). The limitation of the number of such vessels has been 35 and has not changed since 1995.

The total number of pole-and-line vessels showed a continuous declining trend since 1980 (Tables 1-A and B). Suppose vessel size categories 20-49 GRT, 50-199 GRT, and over 200 GRT for 1980-2006 to compare with that for 2006-2023, the number of vessels for each category showed declining trend throughout the period (Table 1-A). The number of vessels both for 50-199 GRT and over 200 GRT showed declining trend throughout the period (Table 1-B). The number of vessels for 50-199 GRT was 34 in 2023 which is 40% of that in 2006. The number of vessels for over 200 GRT showed a declining trend with annual fluctuations, was 20 in 2023, which is 67% of that in 2006.

2. CATCH AND EFFORT TRENDS OF THE MAJOR FISHERIES

2.1. Longline

Longline is classified by the type of license issued by the Government of Japan, i.e., coastal (< 20 GRT and can fish only in Japanese EEZ), small offshore (10-20 GRT), offshore (10-120 GRT), and distant water (> 120 GRT).

Annual distributions of fishing effort of longline in 2022 and 2023 are shown in Fig. 1. In those years, the fishing grounds were in the east-west direction off Japan to Hawaii, the equatorial area between 15°S and 15°N, off Australia and off Peru. The fishing effort of the distant water and offshore longline remained stable at around 200 million hooks in the North Pacific in the 1980s, and then it decreased continuously to 100 million hooks in the early 2000s, and it had further decreased until 2009 (Fig. 2). After 2009, the amount of effort showed a trend of gradual decrease at a level of 35-50 million hooks.

Total catch of four tuna and four billfish species caught by distant water and offshore longline in the North Pacific has been decreased since the highest catch of 119,185 t in 1980 and was 11,734 t in 2023 which is 8% of that in 1980 (Fig. 2). Bigeye has been the dominant species in this fishery in the North Pacific. The bigeye catch, which was stable in the 1980s and about 50,000 t in the late 1980s, showed a declining trend since the 1990s, was less than 10,000 t since 2009, and was less than 5,000 t since 2016. Yellowfin tuna catch ranged between 30,000 t and 50,000 t until the early 1980s. It gradually decreased to less than 5,000 t in 2007. Albacore catch, which fluctuated around 10,000 t until 2001, decreased to about 2,000-6,000 t and kept stable at a low level during the period 2003-2023.

2.2. Purse seine

There are two types of Japanese purse seiners targeting tunas, i.e., single and group purse seine. Historically, a typical group seiner consists of one purse seiner, one searching vessel, and two carrier vessels. Still, the group seiner tended to reduce the number of vessels within each group to reduce costs in recent years. Besides those, coastal purse seiner takes a relatively small number of tunas as a by-catch.

Fishing grounds of Japanese purse seine were widely spread, ranging from 40°N to 10°S, and from 120°E to 180° (Fig. 3). The group seiner operates mainly in the temperate northwestern Pacific. The carrier holds fish in chilled water with ice and unloads those catches. Meanwhile, the single purse seiner (> 349 GRT) operates mainly in the tropical waters of the central and western Pacific, but a part of the vessels seasonally operates in the temperate waters.

The fishing effort was around 9,000 sets in the late 1980s, then decreased to about 6,000 sets in 1998 (Fig 4). The fishing effort generally stayed at about 4,000-6,000 sets in the last decade (Fig. 4). The skipjack catch has been dominant among species in this fishery, followed by yellowfin. The skipjack catch was about 150,000 t until 2008 and then decreased to 80,000 t in 2011 (Fig. 4). After 2011, the skipjack catch showed no clear trend between 80,000 t and 140,000 t (Fig. 4). The statistics in 2023 are provisional, and that skipjack catch is about 65,000 t.

2.3. Pole-and-line

The pole-and-line is composed of three distinct categories, i.e., coastal (< 20 GRT), offshore (10-120 GRT) and distant water (> 120 GRT) vessels in terms of the license of this fishery. Note that some of 19 GRT type vessels obtained offshore licenses since 2007, which are included in offshore category in this document. The pole-and-line can be categorized into large, middle, and small sized vessels which correspond to larger than 300 GRT, 20-300 GRT and less than 20 GRT in vessel size.

Fishing grounds of the pole-and-line were widely spread ranging from 45°N and 10°S, from 120°E to 180°. The fishing ground was rather sequential from north to south and was unlike that in the purse seine fishery. (Fig. 5). The middle-sized vessels generally operate in near shore waters of Japan and their trips are within 10 days. Southernmost fishing area for these vessels, in recent years, is near 15°N, but the important fishing ground is waters north of 25°N, around Japan and adjacent areas (Fig. 5). These vessels primarily fish skipjack and albacore from spring through autumn off the Pacific side of Japan, and harvest relatively small amount of yellowfin and bigeye tuna. They hold fish in cooled water and unload it as fresh fish. The activity of the small pole-and-line vessels is like that of the middle vessels but the area of fishing is limited within the Japanese EEZ, and the trip of these vessels is shorter. On the contrary, the large vessels tend to operate farther off waters from Japan and their trips last for two to three months. Usually, they primarily target for albacore from summer through autumn season in the waters north of 20°N, and skipjack in winter and spring in the waters south of 20°N (Fig. 5). These vessels equip a brine freezer, in which fish caught are immediately stored into a tank filled with cooled brine, and then unloads it as frozen fish.

Generally, fishing effort expressed by fishing days for offshore and distant water pole-and-line rapidly decreased from around 62,000 days in the early 1980s to around 20,000 days in 1994, increased to around 23,000 days in 2000, and then decreased to 6,713 days (preliminary) in 2023 (Fig. 6). Total catch of five tuna species for those fisheries rapidly decreased from around 280,000 t to around 170,000 t during the 1980s, and then gradually decreased from around 130,000 t to 50,000 t until the latest year (Fig. 6). Skipjack is a dominant species for this fishery, but the proportion of skipjack tended to decrease, from 87-78% during 1980-1986 to 87-50% during 2011-2023.

3. RECENT TRENDS FOR MAJOR SPECIES

3.1. Pacific Bluefin Tuna (Table 2-A)

Preliminary total catch of Pacific bluefin tuna (PBF) in 2023 was 9,792 t (Table 2-A), which was similar level with that in previous year (10,112 t). Continuing from 2022, this was the highest level since the strict catch upper limit was implemented in 2015 in accordance with the WCPFC Conservation and Management Measure (CMM). The annual catches of PBF by major fisheries

in 2023 is as follows; purse seine: 4,570 t, troll: 1,160 t, setnet: 1,889 t, longline: 1,556 t, and other fisheries: 593 t.

Japanese longline (LL) fishery had been caught mainly large spawner PBF, which were about 200 cm Fork Length (FL), but the length composition in a recent several years showed multimodal distribution which has the several peaks for less than 200 cm FL (Fig. 7). The peaks at the small spawner (about 150 cm FL or smaller) in the LL composition data became distinctive in 2017 and onward when the availability of young spawner improved in other fisheries as well. Considering with the assessment results, the small spawner shown in the size composition data of longline fishery would be an influx of newly available cohorts, which were protected by the management since 2015, to the fishery rather than the change in migration of spawner fish.

3.2. Albacore (Table 2-B)

The preliminary total catch of albacore in 2023 was 16,274 t, which was smaller than the average of the past five years (2018-2022: 31,963 t). The main Japanese fisheries for albacore are the longline and pole-and-line fisheries. Longline catches have been stable in recent years, at around 12,000 t. Catches by the distant waters pole-and-line fluctuated significantly, ranging from 4,090 t to 36,638 t. This fishery targets both albacore and skipjack, and albacore catches fluctuate depending on the combined catch of both species.

Longline fishery targets the larger fish, while pole-and-line fishery targets much smaller fish (Fig. 8). The size of the albacore caught by the longline in 2023 ranged from 60 to 135 cm. A bimodal size distribution was observed, with peaks at 79 cm and 103 cm FL. The albacore caught by the pole-and-line fishery ranged in size from 46 cm to 88 cm.

3.3. Swordfish (Table 2-C)

Total swordfish catch in 2023 was 5,640 t which is 127.5% of the catch in 2022 (4,424 t). These statistics are preliminary but indicate that the catch somewhat exceeds the average over the past five years (2019-2023: 5,185 t). Swordfish have been caught mainly by offshore and distant water longline, whose catch in 2023 was 3,400 t. The coastal longline catch in 2023 was 1,178 t. Length composition data was collected from longline fishery. The distribution range was approximately from 90 to 220 cm eye-fork length in 2022 and 2023 (Fig. 9).

3.4. Striped Marlin (Table 2-D)

The total striped marlin catch in 2023 was 875 t which is 101.4 % of the catch in 2022 (863 t). These statistics are preliminary and indicate that the catch falls short of the average over the past five years (2019-2023: 1,150 t). In recent years, Japanese pelagic fisheries catch striped marlin as bycatch except for coastal drift-net and part of other longline fisheries that target striped marlin seasonally. Length-composition data was collected from longline fishery. The distribution range was approximately from 100 to 180 cm eye-fork length in 2022 and 2023 (Fig. 10).

3.5. Blue Shark (Table 2-E)

ISC SHARK WG had conducted a benchmark stock assessment of blue shark in 2022. The catch for 2016-2020 was maintained using the annual catch used in the stock assessment (Kai, 2021; Kai and Yano, 2021) and annual catch in 2021-2022 were updated using the almost same estimation method used in the stock assessment in 2021. A total catch of blue shark in 2021 and 2022 was estimated at 9,285 t and 7,986t, respectively. The decreasing trends in total catch since 2017 was mainly due to the decline of catches for longline fisheries

3.6. Shortfin mako (Table 2-F)

ISC SHARK WG had conducted a benchmark stock assessment of shortfin mako shark in 2024. The annual catch for 2017-2022 of shortfin mako was updated using the stock assessment results (Kai, 2023; Kai and Yano, 2023). A total catch of shortfin mako in 2021 and 2022 was estimated at 630 t and 819 t, respectively. The decreasing trends in total catch since 2017 was mainly due to the decline of catches for longline fisheries and large mesh drift-net fishery.

3.7. Others (Bigeye, Skipjack and Yellowfin Tunas) (Table 2-G, H and I)

Preliminary total catch of bigeye in 2023 was 10,728 t which corresponds to 121.4% of the catch in 2022 (8,831 t) and was lower than the average of past five years (2019-2023: 11,2269 t). Total catch of bigeye by Japanese fisheries showed a slight declining trend in the last six years and longline has been the highest proportion among gears in the North Pacific.

Preliminary total catch of skipjack in 2023 was 112,316 t which corresponds to 93.3% of the catch in 2022 (120,385 t) and was lower than the average of past five years (2019-2023: 133,717 t). Most skipjack were caught by pole-and-line and purse seine in the North Pacific Ocean. Total catch of skipjack by Japanese fisheries showed no clear trend in the last six years in the North Pacific Ocean. The lower total catch in 2022 and 2023 were due to lower catch by both pole-and-line and purse seine fisheries.

Preliminary total catch of yellowfin tuna in 2023 was 28,261 t which corresponds to 108.6% of the catch in 2022 (26,018 t) and was lower than the average of past five years (2019-2023: 32,096 t). The yellowfin tuna caught by purse seine has been the highest proportion among gears in the North Pacific Ocean.

4. RESEARCH ACTIVITIES

The Fishery Agency of Japan, in cooperation with the Fisheries Resources Institute (FRI) and local prefectural fisheries experimental stations, has run the nationwide port sampling project for collection of catch, effort and size data of tunas, skipjack, billfishes, and sharks at the major landing ports since the early 1990s. The tagging studies using conventional, archival and pop-up have been conducted by research and training vessels as well as commercial vessels. In addition, there were cooperative works with prefectural fisheries experimental stations and universities. Several cooperative studies were also conducted with foreign countries.

4.1. Pacific Bluefin Tuna

4.1.1. Bluefin Tuna Larvae/Juveniles Research Cruise

Since 2011, larval surveys have been conducted to estimate current main spawning area and period of PBF. In 2023, research cruises were designed to focus on ecological studies of larval/juvenile PBF by R/Vs of FRA, Shunyo-Mar, Yoko-Mar, Hokko-Mar and five prefectural R/Vs. Surveys for larval/juvenile PBF were conducted in the south of Japan around Nansei Islands area, where is a major spawning ground of PBF, from May to July and also in the Sea of Japan, which is another spawning ground of PBF, from July to August. In addition to these two spawning grounds, the survey was conducted in Sanriku-Joban area in the coastal area of northeastern Japan in July and September. In 2023, PBF larvae were captured by all cruises in the spawning grounds. Small juveniles of PBF around 2-5 cm FL were also captured in Nansei Island area and Sanriku-Joban area by small surface-trawl net. In Sanriku-Joban area, we could collect more than 2000 juveniles of tuna species. However, a large part of these fish were Yellowfin and skipjack and only four PBF

juveniles were found in these collected juveniles. Collected samples are being examined by a variety of approaches such as genetic identification, aging, growth analysis, stable isotope, microchemistry and stomach contents analyses to elucidate the survival processes of larval and juvenile PBF in relation to biological and environmental factors, which should help to understand the recruitment mechanism to PBF fisheries around Japan.

4.1.2. Troll survey on age-0 Pacific Bluefin tuna

Recruitment abundance index (standardized CPUE from the Japanese troll fishery) for current PBF stock assessment is based on the sales-slips issued by the commercial markets. To have a better understanding of the recruitment status more timely and precisely than the sales-slip based index, which naturally lacked zero-catch or spatio-temporal information, the FRI commenced using a real-time monitoring survey data of troll fishery's operations in 2011. In this survey, the catch information (Number of fish by species) with its geographic position and Sea Surface Temperature (SST) are sent to the FRI in near-real time fashion using a data transmitter which was equipped on the troll vessels. Furthermore, since 2021 fishing year, IQ-independent charter real-time monitoring surveys were initiated to ensure sufficient operations in each spatial and temporal stratum, in addition to the conventional real-time monitoring.

Based on those data, the FRI reported the latest recruitment index, which informs the relative strength of the cohorts over 2011-to-2023-year classes, to the PBFWG meeting held in Kaohsiung in February 2024 (Fujioka et al., 2024). By function of the data collection system, the PBFWG could see the relative strength of the latest cohort before that cohort turns to age 1 (PBF turns their age in July by the assessment assumption). Totally 3,863 days operational data from 14 real-time monitoring vessels, which targeted for age-0 PBF (i.e., 30-60 cm fork length) during the winter season (November to following February) in the East China Sea were used to standardize the CPUE by Vector Autoregressive Spatio-Temporal (VAST) model formulated a delta-generalized linear mixed model. Estimated indices for 2011-2023 were quite similar to the traditional sales slip index throughout the overlapping period (2011-2016).

In the PBFWG meeting held in February-March 2024, the PBFWG concluded that it was still premature to include this index into the base-case model, however, the WG considered that this index could be useful for qualitative validation of the recruitment estimated by the assessment base case or for the early warning of the sudden recruitment drop. With this regard, the PBF recruitment in 2023 would not be a recruitment failure based on this index.

4.1.3. Tissue Sampling and Technical Development for Close-Kin Analysis for PBF

Tissue sampling for close-kin analysis started in 2015 and around 2,000 individuals were sampled in 2023 as an annual routine. Due to the existence of multiple segregated spawning grounds for PBF stock, the samples only whose spawning ground for adult and hatch ground for juvenile are known were collected. The tissues of large mature adult PBFs (about 120-300 kg in BW; Body Weight) were sampled by coastal longline fishery around Okinawa Islands in late April to early July, while those of young-of-the-year juveniles (about 0.1-0.3 kg in BW) were sampled by troll fishery during summer in the Pacific coastal water off Western Japan, such as Kochi Prefecture. The hatching area of these juveniles can be identified as the water around Nansei archipelago based on the knowledge about the migration pattern of age-0 PBF. For the other spawning ground, the Sea of Japan, young adult PBFs which are either nearly-matured or matured (about 20-60 kg in BW) were sampled in Sakai-minato in June to early July, while those of young-of-the-year juveniles were sampled in Oki islands in late September to early November. All of muscle tissues

sampled were preserved in specific buffer (TNES-Urea 6M buffer) because of the higher stability of content DNA under the room temperature. A part of collected (around 1,500) samples were annually analyzed and sequenced by GRAS-Di technique since 2019.

4.2. Sharks, Billfishes and Swordfish

4.2.1. Port Sampling and the Onboard Research Program in Kesennuma Fishing Port

In 2023, size and sex data of blue shark and shortfin mako were collected from the port sampling in Kesennuma fishing port, located in the northeastern Honshu (the main island of Japan), and the onboard research project for Kesennuma offshore longline fleet throughout the year.

In the port sampling, size data from 11,141 blue shark were collected, and 70% of individuals measured were males. In addition, 69% of males and 70% of females measured were juveniles. In the onboard research program for Kesennuma-offshore longline fleet, the catch number of blue shark was recorded by four size categories (large, middle, small, and extra small). Total of 209,652 blue sharks were recorded and large (processed weight ≥ 15 kg) consisted 48% of all catch with 25% of middle ($11 \text{ kg} \leq \text{processed weight} < 15 \text{ kg}$), 25% of small ($5 \text{ kg} \leq \text{processed weight} < 11 \text{ kg}$) and 3% of extra small (processed weight $< 5 \text{ kg}$).

For shortfin mako, size data from 7,040 individuals was collected in port sampling program, and 52% of individuals measured were males. Among these sharks measured, 783% of males and 100 (99.8)% of females were juveniles. In contrast to blue shark, almost all of sampled female were juvenile in shortfin mako. Total of 8,167 shortfin mako was recorded by size category from the onboard research by Kesennuma-offshore longline fleet. Large (precaudal length $> 200 \text{ cm}$) consisted 3% of all catch with 23% of middle ($150 \text{ cm} < \text{precaudal length} \leq 200 \text{ cm}$), 59% of small ($100 \text{ cm} < \text{precaudal length} \leq 150 \text{ cm}$) and 15% of extra small (precaudal length $\leq 100 \text{ cm}$).

4.2.2. Tagging for Sharks

In 2023, conventional tags were attached to 21 blue sharks in the area around 19 degrees north and 160 degrees east during the research cruise of Japanese research and training vessel (JRTV). The released blue sharks were subadult and adult and the percentage of male was 57%.

4.2.3. Biological Sample Collection

Samples of sagitta, reproductive organ, dorsal fin and anal fin were collected from a total of 434 swordfish, 528 striped marlin, and 174 blue marlin for the collaborative study within ISC billfish working group to estimate biological parameters of billfishes and swordfish. For the study of genetic population structure and other ecological study, muscle tissue was collected from 302 swordfish, 396 striped marlin, and 14 blue marlins.

For sharks, samples of whole body were collected from shortfin mako and salmon shark for the biological study of life history, genetic population structure, and other ecological study. Reproductive organ, muscle, and vertebrae were collected from three adult female shortfin mako to investigate the reproductive cycle, growth, and distribution pattern.

All the samples above were collected by the research cruise (including chartered vessel) and commercial/training longline operation, and sport-fishing conducted in the North Pacific Ocean in 2022.

4.3. Skipjack

4.3.1. Tagging for Skipjack

The FRI has been conducting skipjack tagging research to investigate their migration patterns off Japan. In October 2023, an offshore pole-and-line vessel was chartered to conduct the research. A total of 1,889 skipjack tuna were released, including 80 individuals with archival tags (Lotek, LAT2910). In addition, the FRI has collaborated with Ajinomoto Co., Inc. to conduct skipjack tagging in coastal areas of southwestern Japan. A total of 297 skipjack tuna were released (244 in September 2023 and 53 in March 2024), including 23 individuals with archival tags. Furthermore, three prefectural research and training vessels conducted skipjack tagging in 2023 and 2024. The tagging locations were around Hachijo Island, off Wakayama and Boso area. A total of 164 skipjack tuna were released, including 15 individuals with archival tags and acoustic pingers (AquaSound, AQPX-1040).

4.4. Albacore 鰹

The FRI has been conducting tagging research to investigate female and male albacore distribution and migration in the northwestern Pacific Ocean. In February and March 2024, 4 pop-up tags (Wildlife, miniPAT) and 1 archival tag were attached to albacore in off Wakayama.

5. REFERENCE

- Fujioka K., Asai S., Tsukahara Y., Fukuda H. and Nakatsuka S. (2024) Recruitment abundance index of Pacific bluefin tuna based on real-time troll monitoring survey data using Vector Autoregressive Spatio-Temporal (VAST) model analysis: ISC/24/PBFWG-1/04
- Fujioka K., Asai S., Tsukahara Y., K., Fukuda H. and Nakatsuka S. (2023) Recruitment abundance index of immature Pacific bluefin tuna, derived from real-time monitoring survey data of troll fisheries: ISC/23/PBFWG-1/03
- Kai M. (2021) Update of Japanese annual catches for blue shark caught by Japanese offshore and distant water longliner in the North Pacific Ocean from 1994 to 2020: ISC/21/SHARKWG-2/04
- Kai M. (2023) Update of annual catches for shortfin mako caught by Japanese offshore and distant water longliner in the North Pacific Ocean from 1994 to 2022: ISC/23/SHARKWG-1/05
- Kai, M., Yano, T. (2021) Updated annual catches of blue shark caught by Japanese coastal fisheries in the North Pacific Ocean from 1994 to 2019: ISC/21/SHARKWG-2/05.
- Kai, M., Yano, T. (2023) Updated annual catches of shortfin mako caught by Japanese coastal fisheries in the North Pacific Ocean from 1994 to 2022: ISC/23/SHARKWG-1/04
- MAFF (1982-2008): Gyogyou yousyokugyouseisan toukei nenpou (Yearbook of fisheries and aquaculture production statistics of Japan for 1980-2006), Statistics Department, Minister's Secretariat, Ministry of Agriculture, Forestry and Fishery.
- Tsukahara Y., Asai S., Fukuda H., and Nakatsuka S. (2021) CPUE and Catch at Size for Pacific Bluefin tuna (*Thunnus Orientalis*) caught by Japanese coastal and offshore longline: ISC/21/PBFWG-2/01

6. TABLES

Table 1. Number of Japanese tuna fishing vessels.

A. Number of Japanese tuna fishing vessels operated in the Pacific Ocean by type of fisheries and vessel size based on MAFF (1980-2006).

| Year | Longline fishery ^{*1} | | | | | Total | Purse seine fishery | | | Pole-and-line fishery | | | | | Total |
|------|--------------------------------|-----------|-----------|-------------|----------|-------|--------------------------|----------|-------|------------------------|-----------|-----------|-------------|----------|-------|
| | 1-19 GRT | 20-49 GRT | 50-99 GRT | 100-199 GRT | 200- GRT | | 50-199 GRT ^{*2} | 200- GRT | Total | 1-19 GRT ^{*3} | 20-49 GRT | 50-99 GRT | 100-199 GRT | 200- GRT | |
| 1980 | 821 | 57 | 715 | 103 | 645 | 2,341 | 50 | 16 | 66 | 3,232 | 14 | 350 | 10 | 198 | 3,804 |
| 1981 | 774 | 55 | 706 | 100 | 661 | 2,296 | 50 | 23 | 73 | 3,064 | 10 | 353 | 6 | 179 | 3,612 |
| 1982 | 722 | 43 | 634 | 90 | 589 | 2,078 | 52 | 33 | 85 | 3,011 | 11 | 320 | 6 | 138 | 3,486 |
| 1983 | 561 | 38 | 589 | 93 | 550 | 1,831 | 59 | 36 | 95 | 3,021 | 12 | 297 | 9 | 116 | 3,455 |
| 1984 | 523 | 32 | 538 | 108 | 610 | 1,811 | 54 | 33 | 87 | 2,904 | 8 | 273 | 10 | 105 | 3,300 |
| 1985 | 620 | 28 | 512 | 131 | 628 | 1,919 | 47 | 35 | 82 | 2,754 | 8 | 244 | 9 | 95 | 3,110 |
| 1986 | 536 | 25 | 435 | 168 | 632 | 1,796 | 53 | 38 | 91 | 2,455 | 6 | 224 | 9 | 91 | 2,785 |
| 1987 | 661 | 23 | 348 | 197 | 649 | 1,878 | 47 | 34 | 81 | 2,404 | 6 | 210 | 9 | 89 | 2,718 |
| 1988 | 586 | 21 | 289 | 233 | 649 | 1,778 | 48 | 39 | 87 | 2,613 | 5 | 191 | 11 | 70 | 2,890 |
| 1989 | 650 | 20 | 248 | 238 | 653 | 1,809 | 43 | 37 | 80 | 2,254 | 3 | 187 | 12 | 67 | 2,523 |
| 1990 | 685 | 21 | 227 | 241 | 664 | 1,838 | 43 | 35 | 78 | 2,228 | 4 | 176 | 9 | 66 | 2,483 |
| 1991 | 768 | 19 | 199 | 222 | 682 | 1,890 | 38 | 35 | 73 | 2,277 | 3 | 166 | 10 | 63 | 2,519 |
| 1992 | 793 | 19 | 164 | 206 | 681 | 1,863 | 31 | 38 | 69 | 2,093 | 3 | 156 | 11 | 46 | 2,309 |
| 1993 | 790 | 18 | 138 | 201 | 682 | 1,829 | 27 | 36 | 63 | 1,927 | 3 | 147 | 10 | 43 | 2,130 |
| 1994 | 819 | 21 | 110 | 198 | 675 | 1,823 | 23 | 33 | 56 | 1,830 | 3 | 124 | 10 | 48 | 2,015 |
| 1995 | 738 | 20 | 92 | 187 | 667 | 1,704 | 20 | 31 | 51 | 481 | 3 | 104 | 20 | 46 | 654 |
| 1996 | 711 | 17 | 91 | 155 | 640 | 1,614 | 21 | 32 | 53 | 512 | 3 | 89 | 29 | 43 | 676 |
| 1997 | 698 | 11 | 88 | 145 | 631 | 1,573 | 20 | 35 | 55 | 436 | 2 | 76 | 39 | 45 | 598 |
| 1998 | 712 | 11 | 80 | 129 | 623 | 1,555 | 20 | 35 | 55 | 382 | 2 | 73 | 40 | 46 | 543 |
| 1999 | 703 | 6 | 78 | 119 | 567 | 1,473 | 22 | 36 | 58 | 416 | 1 | 62 | 54 | 46 | 579 |
| 2000 | 732 | 3 | 76 | 111 | 496 | 1,418 | 23 | 37 | 60 | 357 | 1 | 56 | 57 | 47 | 518 |
| 2001 | 777 | 4 | 76 | 110 | 494 | 1,461 | 19 | 36 | 55 | 285 | 1 | 49 | 59 | 47 | 441 |
| 2002 | 780 | 4 | 69 | 110 | 484 | 1,447 | 18 | 36 | 54 | 251 | 1 | 45 | 58 | 48 | 403 |
| 2003 | 764 | 3 | 64 | 99 | 460 | 1,390 | 17 | 36 | 53 | 292 | 1 | 44 | 56 | 44 | 437 |
| 2004 | 702 | 2 | 55 | 77 | 455 | 1,291 | 17 | 36 | 53 | 284 | 1 | 38 | 57 | 43 | 423 |
| 2005 | 694 | 2 | 46 | 59 | 432 | 1,233 | 17 | 36 | 53 | 247 | 1 | 36 | 58 | 45 | 387 |
| 2006 | 709 | 1 | 43 | 54 | 401 | 1,208 | 16 | 36 | 52 | 213 | 1 | 27 | 58 | 36 | 335 |

*1 Longline vessels larger than 50 GRT include those operated in the area other than the Pacific.

*2 50-199 GRT class vessels only include those operated in the Pacific side of northern Japan.

*3 1-19 GRT class vessels before 1995 include those engaged in trolling.

B. Number of Japanese tuna fishing vessels operated in the North Pacific Ocean by type of fisheries and vessel size based on logbook. Values in 2021 and 2022 are provisional.

| Year | Longline fishery | | | | | Purse seine fishery | | | | Pole-and-line fishery | | | |
|------|------------------|--------------|----------------|-------------|-------|-----------------------------|----------------|-------------|-------|-----------------------|---------------|-------------|-------|
| | 10-49 GRT | 50-99 GRT | 100-199 GRT | 200- GRT | Total | 50-199 GRT* ⁴ | 200-499 GRT | 500- GRT | Total | 20-49 GRT | 50-199 GRT | 200- GRT | Total |
| 2006 | 277 | 44 | 52 | 113 | 486 | 31 | 35 | 1 | 67 | 1 | 83 | 30 | 114 |
| 2007 | 279 | 42 | 48 | 89 | 458 | 34 | 36 | 1 | 71 | 1 | 77 | 29 | 107 |
| 2008 | 277 | 42 | 40 | 90 | 449 | 35 | 37 | 1 | 73 | 1 | 69 | 29 | 99 |
| 2009 | 277 | 38 | 33 | 81 | 429 | 33 | 36 | 3 | 72 | 1 | 68 | 28 | 97 |
| 2010 | 290 | 29 | 28 | 98 | 445 | 31 | 35 | 4 | 70 | 1 | 66 | 28 | 95 |
| 2011 | 273 | 24 | 25 | 99 | 421 | 33 | 36 | 4 | 73 | 0 | 63 | 28 | 91 |
| 2012 | 265 | 21 | 21 | 92 | 399 | 34 | 37 | 4 | 75 | 0 | 60 | 27 | 87 |
| 2013 | 260 | 20 | 23 | 87 | 390 | 34 | 37 | 4 | 75 | 0 | 55 | 25 | 80 |
| 2014 | 250 | 18 | 21 | 90 | 379 | 33 | 37 | 3 | 73 | 1 | 54 | 25 | 80 |
| 2015 | 239 | 18 | 24 | 80 | 361 | 30 | 35 | 5 | 70 | 1 | 51 | 24 | 76 |
| 2016 | 234 | 16 | 16 | 64 | 330 | 38 | 33 | 4 | 75 | 1 | 50 | 25 | 76 |
| 2017 | 233 | 15 | 15 | 50 | 313 | 37 | 34 | 4 | 75 | 1 | 48 | 31 | 80 |
| 2018 | 230 | 14 | 16 | 63 | 323 | 35 | 30 | 4 | 69 | 1 | 44 | 25 | 70 |
| 2019 | 230 | 13 | 17 | 51 | 311 | 35 | 31 | 5 | 71 | 1 | 42 | 24 | 67 |
| 2020 | 228 | 11 | 15 | 42 | 296 | 34 | 31 | 6 | 71 | 1 | 38 | 22 | 61 |
| 2021 | 201 | 10 | 17 | 48 | 276 | 32 | 29 | 7 | 68 | 1 | 36 | 22 | 59 |
| 2022 | 214 | 6 | 16 | 41 | 277 | 22 | 31 | 8 | 61 | 1 | 33 | 22 | 56 |
| 2023 | 195 | 6 | 13 | 38 | 252 | 19 | 25 | 9 | 53 | 1 | 34 | 20 | 55 |

*4 50-199 GRT class vessels only include those operated in the Pacific side of northern Japan.

Table 2. Catch in weight (t) by species by fisheries in the North Pacific.**A. Pacific bluefin tuna**

| Year | Purse Seine | | Dist. & Off. Longline ^{*5} | | Coastal Longline | Troll | Pole-and-line | Set-net | Others ^{*7} | Total |
|--------------------|-------------|----------|-------------------------------------|---------------|------------------|-------|---------------|---------|----------------------|--------|
| | Tuna PS | Small PS | Noth Pcific | South Pacific | | | | | | |
| 2017 | 3,341 | 1,199 | 21 | 6 | 892 | 605 | 49 | 2,221 | 665 | 9,000 |
| 2018 | 3,225 | 825 | 21 | 0 | 679 | 371 | 9 | 645 | 431 | 6,205 |
| 2019 | 3,213 | 1,251 | 25 | 0 | 977 | 720 | 0 | 951 | 372 | 7,509 |
| 2020 | 3,208 | 752 | 75 | 0 | 1,341 | 760 | 1 | 1,342 | 532 | 8,011 |
| 2021 | 3,185 | 1,014 | 80 | 0 | 1,432 | 653 | 0 | 1,742 | 512 | 8,617 |
| 2022 | 3,671 | 1,031 | 80 | 0 | 1,519 | 1,079 | 13 | 2,126 | 593 | 10,112 |
| 2023 ^{*6} | 3,800 | 770 | 80 | 0 | 1,477 | 1,160 | 24 | 1,889 | 593 | 9,792 |

*5: Distant-water and Offshore longline vessels are mainly 20 GRT or larger, and most of coastal longline vessels are smaller than 20 GRT.

*6: Most recent year's catch value is provisional.

*7: Others include drift net, handline, trawl, other longline, and unclassified fisheries.

B. Albacore

| Year | Longline | | Pole-and-line | | | | | Troll | Set-net | Others | Total |
|------|----------------------------|---------|----------------|----------|---------|-----------|-------------|-------|---------|--------|----------|
| | Distant Water + Offshore*8 | Coastal | Distant Waters | Offshore | Coastal | Drift-net | Purse seine | | | | |
| 2018 | 3,071 | 10,121 | 9,362 | 8,394 | 119 | 35 | 3,039 | 78 | 13 | 70 | 34,302 |
| 2019 | 2,841 | 9,375 | 4,669 | 3,662 | 177 | 9 | 1,045 | 543 | 27 | 95 | 22,443 |
| 2020 | 2,415 | 10,241 | 23,802 | 12,578 | 254 | 7 | 5,961 | 784 | 25 | 159 | 56,226 |
| 2021 | 4,157 | 14,512 | 6,869 | 4,043 | 224 | 3 | 92 | 428 | 11 | 232 | 30,571 |
| 2022 | 2,756 | 8,278 | 2,234 | 1,770 | 86 | 31 | 726 | 216 | 18 | 159 | 16,274 |
| 2023 | (2,756) | (8,278) | (2,234) | (1,770) | (86) | (31) | (726) | (216) | (18) | (159) | (16,274) |

*8 Category Distant Water + Offshore LL includes training/research vessels

() different data source or carry over from previous year

C. Swordfish

| Year | Longline | | | Drift-net | Bait fishing | Net fishing | Trap-net | Others | Total |
|--------------------|--------------------------|---------|--------|-----------|--------------|-------------|----------|--------|-------|
| | Distant Water + Offshore | Coastal | Others | | | | | | |
| 2018 | 3,212 | 1,801 | 2 | 230 | 267 | NA | 5 | 480 | 5,997 |
| 2019 | 2,601 | 1,307 | 2 | 242 | 210 | 0 | 6 | 339 | 4,706 |
| 2020 | 3,842 | 1,471 | 4 | 290 | 305 | 0 | 7 | 179 | 6,098 |
| 2021 | 2,985 | 1,238 | 8 | 301 | 251 | 0 | 4 | 270 | 5,057 |
| 2022 ^{*9} | 2,260 | 1,102 | 4 | 459 | 283 | 0 | 4 | 313 | 4,424 |
| 2023 ^{*9} | 3,400 | 1,178 | 4 | 459 | 283 | 0 | 4 | 313 | 5,640 |

*9 Catches for 2022 and 2023 are preliminary.

D. Striped Marlin

| Year | Longline | | | | Drift-net | Bait fishing | Net fishing | Trap-net | Others | Total |
|---------------------|--------------------------|---------|--------|-----|-----------|--------------|-------------|----------|--------|-------|
| | Distant Water + Offshore | Coastal | Others | | | | | | | |
| 2018 | 147 | 711 | 28 | 278 | 36 | NA | 28 | 52 | 1,280 | |
| 2019 | 222 | 889 | 29 | 241 | 39 | NA | 29 | 61 | 1,510 | |
| 2020 | 196 | 896 | 49 | 155 | 25 | 0 | 37 | 32 | 1,390 | |
| 2021 | 184 | 708 | 17 | 95 | 17 | NA | 31 | 60 | 1,112 | |
| 2022 ^{*10} | 137 | 451 | 15 | 138 | 23 | NA | 27 | 71 | 863 | |
| 2023 ^{*10} | 133 | 468 | 15 | 138 | 23 | NA | 27 | 71 | 875 | |

*10 Catches for 2022 and 2023 are preliminary.

E. Blue shark

| Year | Longline | | | | Large mesh driftnet | Bait fishing | Trapnet | Others | Total |
|------|---------------|----------|---------|--------|---------------------|--------------|---------|--------|--------|
| | Distant Water | Offshore | Coastal | Others | | | | | |
| 2017 | 4,387 | 4,853 | 343 | 212 | 1,366 | 1 | 4 | 0 | 11,166 |
| 2018 | 4,081 | 4,608 | 263 | 159 | 1,236 | 1 | 40 | 0 | 10,388 |
| 2019 | 3,726 | 4,351 | 209 | 162 | 1,149 | 1 | 35 | 0 | 9,634 |
| 2020 | 3,134 | 3,540 | 213 | 185 | 1,119 | 2 | 59 | 1 | 8,252 |
| 2021 | 3,655 | 3,471 | 416 | 232 | 1,484 | 1 | 25 | 1 | 9,285 |
| 2022 | 3,685 | 2,396 | 605 | 207 | 1,062 | 1 | 29 | 0 | 7,986 |

F. Shortfin mako

| Year | Longline | | | Large mesh driftnet | Trapnet and others | Total |
|------|--|---------------------------------------|-------------------|---------------------|--------------------|-------|
| | Offshore and Distant-water (Shallow-set) | Offshore and Distant-water (Deep-set) | Coastal and other | | | |
| 2017 | 777 | 74 | 23 | 271 | 10 | 1,155 |
| 2018 | 893 | 83 | 19 | 223 | 28 | 1,247 |
| 2019 | 785 | 83 | 15 | 214 | 3 | 1,100 |
| 2020 | 488 | 57 | 4 | 194 | 16 | 759 |
| 2021 | 406 | 51 | 16 | 133 | 23 | 630 |
| 2022 | 589 | 23 | 6 | 161 | 41 | 819 |

G. Bigeye

| Year | Longline | Pole-and-line | Purse seine | Gillnet | Set-net | Troll | Other | Total |
|------|----------|---------------|-------------|---------|---------|-------|-------|--------|
| 2018 | 11,630 | 1,432 | 3,471 | 1 | 0 | 80 | 84 | 16,698 |
| 2019 | 10,693 | 548 | 1,444 | 1 | 0 | 110 | 113 | 12,909 |
| 2020 | 9,158 | 1,125 | 1,622 | 0 | 1 | 69 | 135 | 12,110 |
| 2021 | 8,268 | 1,576 | 1,548 | 0 | 3 | 78 | 81 | 11,554 |
| 2022 | 6,360 | 1,363 | 927 | 0 | 1 | 80 | 100 | 8,831 |
| 2023 | 7,568 | 1,345 | 1,634 | 0 | 1 | 80 | 100 | 10,728 |

H. Skipjack

| Year | Longline | Pole-and-line | Purse seine | Gillnet | Set-net | Troll | Other | Total |
|------|----------|---------------|-------------|---------|---------|-------|-------|---------|
| 2018 | 21 | 78,998 | 125,119 | 91 | 494 | 1,154 | 133 | 206,010 |
| 2019 | 38 | 73,592 | 84,054 | 96 | 246 | 1,387 | 110 | 159,523 |
| 2020 | 24 | 48,804 | 55,352 | 70 | 335 | 949 | 86 | 105,620 |
| 2021 | 70 | 84,837 | 82,805 | 144 | 580 | 2,161 | 148 | 170,745 |
| 2022 | 45 | 53,236 | 65,789 | 125 | 219 | 900 | 71 | 120,385 |
| 2023 | 41 | 44,235 | 66,725 | 125 | 219 | 900 | 71 | 112,316 |

I. Yellowfin tuna

| Year | Longline | Pole-and-line | Purse seine | Gillnet | Set-net | Troll | Other | Total |
|------|----------|---------------|-------------|---------|---------|-------|-------|--------|
| 2018 | 7,955 | 3,519 | 38,868 | 6 | 77 | 1,738 | 587 | 52,750 |
| 2019 | 9,901 | 2,930 | 27,039 | 4 | 208 | 2,070 | 778 | 42,930 |
| 2020 | 6,382 | 3,070 | 18,224 | 13 | 125 | 2,008 | 846 | 30,668 |
| 2021 | 7,766 | 3,717 | 17,980 | 7 | 206 | 2,160 | 767 | 32,603 |
| 2022 | 6,246 | 2,520 | 13,872 | 4 | 378 | 2,180 | 818 | 26,018 |
| 2023 | 7,081 | 2,554 | 15,246 | 4 | 378 | 2,180 | 818 | 28,261 |

FIGURES

Longline fishery




Fig 1. Distribution of fishing effort (Number of hooks) for the Japanese distant water and offshore longline fisheries in the Pacific, 2022-2023.




Fig 2. Historical catches in weight (t) for major species and fishing effort (Number of hooks in million) of the Japanese distant water and offshore longline fisheries (not including small offshore fishery) in the North Pacific. PBF: Pacific bluefin tuna, ALB: albacore, BET: bigeye, YFT: yellowfin tuna, SWO: swordfish, MLS: striped marlin, BUM: blue marlin, BLM: black marlin. The black solid line indicates number of hooks. Values in 2022 and 2023 are provisional.

Purse Seine Fishery




Fig 3. Distribution of fishing effort (number of sets) for the Japanese purse seine fishery in the Pacific, 2022-2023.




Fig 4. Historical catches in weight (t) for major species and fishing efforts (number of sets) of the Japanese purse seine fishery in the North Pacific. SKJ: skipjack, YFT: yellowfin tuna, BET: bigeye, PBF: Pacific bluefin tuna, ALB: albacore. Since 2011, Japanese logbook data has included records of purse seine operations that do not specifically target tunas. The black solid line represents the overall number of sets by target and non-target tunas. The black dashed line denotes the total number of sets by targeting vessels since 2011. The statistics for 2023 are still provisional.

Pole-and-Line Fishery




Fig 5. Distribution of fishing effort (number of days) of the Japanese pole-and-line fishery (larger than 20 GRT vessels) in the Pacific, 2022-2023. Distribution of fishing effort in 2023 is provisional (right panel).




Fig.6. Historical catches in weight (t) for major species and fishing effort (Number of fishing days) of Japanese distant water and offshore fisheries in the North Pacific. SKJ: skipjack, ALB: albacore, YFT: yellowfin tuna, PBF: Pacific bluefin tuna, BET: bigeye. The catch for PBF includes the catch by coastal pole-and-line (less than 20 GRT vessels) fishery. Value in 2023 is provisional.




Fig.7. Annual relative length frequency distribution (simply summing up all measurements) for Pacific bluefin tuna (PBF) caught by longline in 2022 (left) and 2023 (right). Texts in each graph indicate gear, species, year, and the number of fish measured.




Fig.8. Annual relative length frequency distribution (simply summing up all measurements) for albacore (ALB) caught by longline (upper two panels) and pole-and-line (lower two panels) in 2022 (left) and 2023 (right). Texts in each graph indicate gear, species, year, and the number of fish measured.




Fig.9. Annual relative length frequency distribution (simply summing up all measurements) for swordfish (SWO) caught by longline in 2022 (left) and 2023 (right). Texts in each graph indicate gear, species, year, and the number of fish measured.




Fig.10. Annual relative length frequency distribution (simply summing up all measurements) for striped marlin (MLS) caught by longline in 2022 (left) and 2023 (right). Texts in each graph indicate gear, species, year, and the number of fish measured.